



Original article

An exponential Chebyshev second kind approximation for solving high-order ordinary differential equations in unbounded domains, with application to Dawson's integral



Mohamed A. Ramadan^a, Kamal R. Raslan^b, Talaat S. El Danaf^c, Mohamed A. Abd El Salam^{b,*}

^a Mathematics Department, Faculty of Science, Menoufia University, Shebin El-Koom, Egypt

^b Mathematics Department, Faculty of Science, Al-Azhar University, Nasr-City, Cairo, Egypt

^c Department of Mathematics and Statistics, Taibah University Madinah Munawwarah, Saudi Arabia

ARTICLE INFO

Article history:

Received 17 May 2016

Revised 21 June 2016

Accepted 3 July 2016

Available online 18 July 2016

Keywords:

Exponential second kind Chebyshev functions

High-order ordinary differential equations

Collocation method

Dawson's integral

ABSTRACT

A new exponential Chebyshev operational matrix of derivatives based on Chebyshev polynomials of second kind (ESC) is investigated. The new operational matrix of derivatives of the ESC functions is derived and introduced for solving high-order linear ordinary differential equations with variable coefficients in unbounded domain using the collocation method. As an application the introduced method is used to evaluate Dawson's integral by solving its differential equation. The corresponding differential equation to Dawson's integral is a boundary value problem with conditions tends to infinity. The obtained numerical results are compared with the exact solution and showed good accuracy.

© 2016 Egyptian Mathematical Society. Production and hosting by Elsevier B.V.

This is an open access article under the CC BY-NC-ND license.

(<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

1. Introduction

Spectral methods have been developed rapidly through the last years for the numerical solutions of differential equations. Compared to other numerical methods, spectral methods give high accuracy and have wide range of applications in many mathematical problems and physical phenomena. The main idea of spectral methods is to approximate the solutions of differential equations by means of truncated series of some orthogonal polynomials. The most common spectral methods used to solve ordinary differential equations (ODEs) are tau, collocation, and Galerkin methods. Siyyam [1] used Laguerre tau method to solve ODEs while Parand and Razzaghi [2] used the same method with the same equations but with rational Legendre as the basis function. Guo et al. [3] and Wang et al. [4] employed the Legendre collocation method to solve the initial value problems and Awoyemi and Idowu [5] used the hybrid collocation with third order ODEs. Galerkin method is also applied for solving ODEs [6,7]. Doha et al. used the generalized Jacobi polynomials for solving ODEs [8–11].

Chebyshev polynomials are one of the most important orthogonal polynomials, which are widely used with spectral methods [12].

The Chebyshev first kind $T_n(x)$ are orthogonal polynomials on the finite interval $[-1, 1]$, these polynomials have many applications in numerical analysis [12], and numerous studies show the merits of them in various applications in fluid mechanics. One of the applications of Chebyshev polynomials is the solution of ODEs with initial and boundary conditions, with collocation points [13,14]. Many studies are considered on the finite interval $[0, 1]$ with the help of usual transformation maps the Chebyshev to the shifted Chebyshev polynomial. Therefore, under a transformation that maps the interval $[-1, 1]$ into a semi-infinite domain $[0, \infty)$, several research groups successfully applied spectral methods to solve differential equations [15–26], their transformation maps the Chebyshev polynomials to the rational Chebyshev functions (RC) and defined by.

$$R_n(x) = T_n\left(\frac{x-1}{x+1}\right). \quad (1)$$

Furthermore, Koc and Kurnaz [27] have proposed a modified type of Chebyshev polynomials as an alternative to the solutions of the partial differential equations defined in real domain. In their study, the basis functions called exponential Chebyshev (EC) functions $E_n(x)$ which are orthogonal in $(-\infty, \infty)$. This kind of

* Corresponding author.

E-mail addresses: ramadanmohamed13@yahoo.com (M.A. Ramadan), kamal_raslan@yahoo.com (K.R. Raslan), talaat11@yahoo.com (T.S. El Danaf), mohamed_salam1985@yahoo.com (M.A. Abd El Salam).

extension tackles the problems over the whole real domain. The EC functions are defined as

$$E_n(x) = T_n\left(\frac{e^x - 1}{e^x + 1}\right). \tag{2}$$

In our previous report [28] we introduced a modified form of the operational matrix of the derivatives by processing the truncation made by Koc and Kurnaz [27] and applied it to ODEs defined in whole rang. Recently, we reported a new operational matrix of derivatives of EC functions for solving ODEs in unbounded domains [29].

In this paper we introduce a new operational matrix of derivatives based on exponential Chebyshev of the second kind (ESC) functions and employ it to solve ODEs with variable coefficients in unbounded domains using the collocation method.

As an application of our method we find approximate solution to Dawson’s integral by solving its differential equation with the subjected condition that tends to infinity. The high-order linear nonhomogeneous differential equations that considered here in this paper is

$$\sum_{k=0}^m q_k(x) \phi^{(k)}(x) = f(x), \quad -\infty < x < \infty, \tag{3}$$

with the mixed conditions

$$\sum_{k=0}^{m-1} \sum_{j=0}^J d_{ij}^k \phi^{(k)}(b_j) = \alpha_i, \tag{4}$$

$$-\infty < b_j < \infty, \quad i = 0, 1, \dots, m - 1, \quad j = 0, 1, \dots, J$$

where, $q_k(x)$ and $f(x)$ are continuous functions on the interval $(-\infty, \infty)$, d_{ij}^k, b_j and α_i are appropriate constants, or b_j may tends to $\pm \infty$ (the boundary condition tends to infinity).

2. The exponential Chebyshev functions of second kind

In this section we list the definition and some properties of the ESC functions.

2.1. Definition of ESC functions

The ESC function of the form

$$E_n^U(x) = U_n\left(\frac{e^x - 1}{e^x + 1}\right), \tag{5}$$

where $U_n(x)$ is the Chebyshev polynomials of the second kind which are orthogonal polynomials of degree n in x defined on the interval $[-1, 1]$ (see Ref. [12] and [30] for more details).

And the corresponding recurrence relation takes the following form

$$E_0^U(x) = 1, \quad E_1^U(x) = 2\left(\frac{e^x - 1}{e^x + 1}\right),$$

$$E_{n+1}^U(x) = 2\left(\frac{e^x - 1}{e^x + 1}\right)E_n^U(x) - E_{n-1}^U(x). \quad n \geq 1 \tag{6}$$

2.2. ESC functions are orthogonal

The ESC functions are orthogonal in the interval $(-\infty, \infty)$ with respect to the weight function $w(x)$ which is given by $4e^{3x/2}(e^x + 1)^{-3}$, with the orthogonality condition

$$\langle E_n^U(x), E_m^U(x) \rangle = \int_{-\infty}^{\infty} E_n^U(x)E_m^U(x)w(x)dx = \frac{\pi}{2}\delta_{nm}, \tag{7}$$

where, δ_{nm} is the Kronecker delta function and $\langle *, * \rangle$ is the inner product notation.

Also the product relation of ESC functions is given by

$$\left(\frac{e^x - 1}{e^x + 1}\right)E_n(x) = \frac{1}{2}[E_{n+1}^U(x) + E_{n-1}^U(x)] \tag{8}$$

2.3. Function expansion in terms of ESC functions

A function $h(x)$ is well defined over the interval $(-\infty, \infty)$ and can be expanded in terms of ESC functions as

$$h(x) = \sum_{i=0}^{\infty} a_i E_i^U(x), \tag{9}$$

where

$$a_i = \frac{2}{\pi} \int_{-\infty}^{\infty} E_i^U(x)h(x)w(x)dx.$$

If the summation in expression (9) is truncated to N where $N < \infty$ it takes the following form

$$h(x) \cong \sum_{i=0}^N a_i E_i^U(x), \tag{10}$$

also, the (k) th-order derivative of $h(x)$ can be written as

$$h^{(k)}(x) \cong \sum_{i=0}^N a_i (E_i^U(x))^{(k)} \tag{11}$$

where $(E_n^U(x))^{(0)} = E_n^U(x)$.

2.4. The operational matrix

The new representation of ESC functions is presented as follows.

The Chebyshev polynomials of first kind $T_n(x)$ can be expressed in terms of x^n in different formulas found in Ref. [12], one of them is

$$T_n(x) = \sum_{k=0}^{[n/2]} (-1)^k 2^{n-2k-1} \frac{n}{n-k} \binom{n-k}{k} x^{n-2k}, \quad 2k \leq n. \tag{12}$$

Similar relation found in [30,31] for the Chebyshev polynomials of second kind $U_n(x)$ takes the following form

$$U_n(x) = \sum_{k=0}^{[n/2]} (-1)^k \binom{n-k}{k} (2x)^{n-2k}, \tag{13}$$

by the help of properties of gamma function the previous relation takes the form

$$U_n(x) = \sum_{k=0}^{[n/2]} (-1)^k 2^{n-2k} \frac{\Gamma(n-k+1)}{\Gamma(k+1)\Gamma(n-2k+1)} x^{n-2k}, \quad n > 0, \tag{14}$$

where, $[\frac{n}{2}]$ denotes the integer part of the value $\frac{n}{2}$.

If we use the expression $v(x) = \frac{e^x - 1}{e^x + 1}$ in the ESC functions, we can express it explicitly in terms of powers of $v(x)$ as

$$E_n^U(x) = \sum_{k=0}^{[n/2]} (-1)^k 2^{n-2k} \binom{n-k}{k} (v(x))^{n-2k}, \tag{15}$$

from previous relation with simple modification we can define:

if n is even number

$$E_n^U(x) = E_{2l}^U(x) = \sum_{j=0}^l (-1)^{l-j} 2^{2j} \binom{l+j}{l-j} (v(x))^{2j}, \tag{16}$$

By substituting the collocation points x_i in (22), we have the system

$$[\phi^{(k)}(x_i)] = V^{(k)}(x_i)M^T A, \quad i = 0, 1, \dots, N$$

or

$$\Phi^{(k)} = \begin{bmatrix} \phi^{(k)}(x_0) \\ \phi^{(k)}(x_1) \\ \vdots \\ \phi^{(k)}(x_N) \end{bmatrix} = V^{(k)}M^T A, \tag{27}$$

where

$$V^{(k)} = \begin{bmatrix} V^{(k)}(x_0) \\ V^{(k)}(x_1) \\ \vdots \\ V^{(k)}(x_N) \end{bmatrix} = \begin{bmatrix} (v^0(x))_{x=x_0}^{(k)} & (v^1(x))_{x=x_0}^{(k)} & \dots & (v^N(x))_{x=x_0}^{(k)} \\ (v^0(x))_{x=x_1}^{(k)} & (v^1(x))_{x=x_1}^{(k)} & \dots & (v^N(x))_{x=x_1}^{(k)} \\ \vdots & \vdots & \ddots & \vdots \\ (v^0(x))_{x=x_N}^{(k)} & (v^1(x))_{x=x_N}^{(k)} & \dots & (v^N(x))_{x=x_N}^{(k)} \end{bmatrix}.$$

Consequently, from the matrices form (26) and (27), we obtain the fundamental matrix equation for Eq. (3) in the following form

$$\sum_{k=0}^m Q_k V^{(k)} M^T A = F, \tag{28}$$

next, we can obtain the corresponding matrices form for the conditions (4) as follows, using the relation (19) with same procedures for $x = b_j$ we have the fundamental matrix equation corresponding to the mixed conditions (4) as

$$\sum_{k=0}^{m-1} \sum_{j=0}^J d_{ij}^k V^{(k)}(b_j) M^T A = [\alpha_i], \tag{29}$$

$$-\infty < b_j < \infty, i = 0, 1, \dots, m - 1, j = 0, 1, \dots, J$$

4. Description of the collocation method

The fundamental matrix (28) for Eq. (3) corresponding to system of $(N + 1)$ algebraic equations for the $(N + 1)$ unknown coefficients a_0, a_1, \dots, a_N .

We can write Eq. (28) shortly as

$$WA = F \text{ or } [W; F], \tag{30}$$

so that

$$W = (w_{ij}) = \sum_{k=0}^m Q_k V^{(k)} M^T, \quad i, j = 0, 1, \dots, N$$

we can obtain the matrix form for the mixed conditions by means of (29) as

$$R_i A = [\alpha_i] \text{ or } [R_i; \alpha_i], i = 0, 1, \dots, m - 1 \tag{31}$$

where

$$R_i = (r_{i,j}) = \sum_{k=0}^{m-1} \sum_{j=0}^J d_{ij}^k V^{(k)}(b_j) M^T.$$

To obtain the solution of Eq. (3) under the conditions (4), we replace the rows of matrices (31) by any m rows of the matrix (30).

Then, we have the required augmented matrix as

$$[W^*; F^*] = \begin{bmatrix} W_{00} & W_{01} & \dots & W_{0N} & ; & f(x_0) \\ W_{10} & W_{11} & \dots & W_{1N} & ; & f(x_1) \\ \dots & \dots & \dots & \dots & ; & \dots \\ W_{N-m,0} & W_{N-m,1} & \dots & W_{N-m,N} & ; & f(x_{N-m}) \\ r_{00} & r_{01} & \dots & r_{0N} & ; & \alpha_0 \\ r_{10} & r_{11} & \dots & r_{1N} & ; & \alpha_1 \\ \dots & \dots & \dots & \dots & ; & \dots \\ r_{m-1,0} & r_{m-1,1} & \dots & r_{m-1,N} & ; & \alpha_{m-1} \end{bmatrix}, \tag{32}$$

or the corresponding matrix equation

$$W^* A = F^*,$$

we always found that the rank $(W^*) = \text{rank } [W^*; F^*] = N + 1$, it means that the matrix inverse of W^* can be obtained, then, we can write

$$A = (W^*)^{-1} F^*. \tag{33}$$

Thus, the coefficient $a_i, i = 0, 1, \dots, N$ are uniquely determined by Eq. (33), and the approximate solution of the given differential equation obtained from Eq. (20).

5. Illustrative examples

Now, we take some test examples to examine our method, the numerical computations are carried out by the MATHEMATICA.7.0 software program. In the rest of the paper the appendix section contains “code form” made for Example 5.1 and 5.4 with results.

Example 5.1. Consider the following second order non-homogeneous boundary value problem with variable coefficients [28]

$$\phi'' - \frac{1}{1 + e^x} \phi' - \frac{15e^{2x}}{(1 + e^x)^2} \phi = \frac{e^{2x}}{(1 + e^x)^6}, \quad x \in (-\infty, \infty) \tag{34}$$

where, the boundary conditions are

$$\phi(x) = 0 \text{ when } x \rightarrow \infty \text{ and } \phi(x) = 1 \text{ when } x \rightarrow -\infty.$$

The fundamental matrix for the pervious equation is

$$\left\{ Q_0 V^{(0)}(M^T)^0 + Q_1 V^{(1)}(M^T)^1 + Q_2 V^{(2)}(M^T)^2 \right\} A = F,$$

and

$$q_0 = \frac{-15e^{2x}}{(1 + e^x)^2}, \quad q_1 = \frac{-1}{(1 + e^x)}, \quad q_2 = 1, \quad f(x) = \frac{e^{2x}}{(1 + e^x)^6},$$

for $N = 4$, the collocation points are

$$x_0 \rightarrow \infty, \quad x_1 = \ln(3 + 2\sqrt{2}), \quad x_2 = 0,$$

$$x_3 = \ln(3 - 2\sqrt{2}), \quad x_4 \rightarrow -\infty.$$

And, it is clear that Q_2 is the identity matrix where, matrices $V^{(0)}, V^{(1)}, V^{(2)}, Q_1, Q_0$ and M are in the following form

$$Q_0 = \begin{bmatrix} -15 & 0 & 0 & 0 & 0 \\ 0 & \frac{-15}{8}(3 + 2\sqrt{2}) & 0 & 0 & 0 \\ 0 & 0 & \frac{-15}{4} & 0 & 0 \\ 0 & 0 & 0 & \frac{15}{8}(-3 + 2\sqrt{2}) & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix},$$

$$M = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 2 & 0 & 0 & 0 \\ -1 & 0 & 4 & 0 & 0 \\ 0 & -4 & 0 & 8 & 0 \\ 1 & 0 & -12 & 0 & 16 \end{bmatrix},$$

Table 1
Comparing the approximate and exact solutions for Example 5.2.

x	Exact solution	EC with $N = 10$	Absolute error	EC with $N = 16$	Absolute error
-3.0	0.0993279	0.0992278	$1.0 \times e^{-4}$	0.0993274	$5.13535 \times e^{-7}$
-2.5	0.163071	0.163049	$2.18 \times e^{-5}$	0.163071	$1.35492 \times e^{-7}$
-2.0	0.265802	0.265983	$1.8 \times e^{-4}$	0.265803	$7.49985 \times e^{-7}$
-1.5	0.425096	0.425182	$8.63 \times e^{-5}$	0.425095	$1.00225 \times e^{-6}$
-1.0	0.648054	0.647794	$2.6 \times e^{-4}$	0.648055	$1.0313 \times e^{-6}$
-0.5	0.886819	0.886899	$7.98 \times e^{-5}$	0.886818	$7.33901 \times e^{-7}$
0.0	1	1.0002	$2.02 \times e^{-4}$	1.0	$4.2298 \times e^{-7}$
0.5	0.886819	0.886899	$7.98 \times e^{-5}$	0.886818	$7.33901 \times e^{-7}$
1.0	0.648054	0.647794	$2.6 \times e^{-4}$	0.648055	$1.0313 \times e^{-6}$
1.5	0.425096	0.425182	$8.63 \times e^{-5}$	0.425095	$1.00225 \times e^{-6}$
2.0	0.265802	0.265983	$1.8 \times e^{-4}$	0.265803	$7.49985 \times e^{-7}$
2.5	0.163071	0.163049	$2.18 \times e^{-5}$	0.163071	$1.35492 \times e^{-7}$
3.0	0.0993279	0.0992278	$1.0 \times e^{-4}$	0.0993274	$5.13535 \times e^{-7}$

Table 2
The L_2, L_∞ error norms for Example 5.2.

	L_2	L_∞
$N = 10$	$4.354 \times e^{-7}$	$2.60092 \times e^{-4}$
$N = 12$	$1.00329 \times e^{-8}$	$3.75879 \times e^{-5}$
$N = 16$	$6.89304 \times e^{-12}$	$1.0313 \times e^{-6}$

$$Q_1 = \begin{bmatrix} 0 & 0 & 0 & 0 & 0 \\ 0 & \frac{1}{4}(-2 + \sqrt{2}) & 0 & 0 & 0 \\ 0 & 0 & -\frac{1}{2} & 0 & 0 \\ 0 & 0 & 0 & \frac{1}{4}(-2 - \sqrt{2}) & 0 \\ 0 & 0 & 0 & 0 & -1 \end{bmatrix},$$

$$V^{(0)} = \begin{bmatrix} 1 & 1 & 1 & 1 & 1 \\ 1 & \frac{1}{\sqrt{2}} & \frac{1}{2} & \frac{1}{2\sqrt{2}} & \frac{1}{4} \\ 1 & 0 & 0 & 0 & 0 \\ 1 & \frac{-1}{\sqrt{2}} & \frac{1}{2} & \frac{-1}{2\sqrt{2}} & \frac{1}{4} \\ 1 & -1 & 1 & -1 & 1 \end{bmatrix},$$

$$V^{(1)} = \begin{bmatrix} 0 & 0 & 0 & 0 & 0 \\ 0 & \frac{1}{4} & \frac{1}{2\sqrt{2}} & \frac{3}{8} & \frac{1}{2\sqrt{2}} \\ 0 & \frac{1}{2} & 0 & 0 & 0 \\ 0 & \frac{1}{4} & \frac{-1}{2\sqrt{2}} & \frac{3}{8} & \frac{-1}{2\sqrt{2}} \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix},$$

$$V^{(2)} = \begin{bmatrix} 0 & 0 & 0 & 0 & 0 \\ 0 & \frac{-1}{4\sqrt{2}} & \frac{-1}{8} & 0 & \frac{1}{8} \\ 0 & 0 & \frac{1}{2} & 0 & 0 \\ 0 & \frac{1}{4\sqrt{2}} & \frac{-1}{8} & 0 & \frac{1}{8} \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix},$$

and the augmented matrix for the boundary conditions with $N = 4$, is: for $x \rightarrow \infty, \phi = 0$ give $[1 \ 2 \ 3 \ 4 \ 5 \ ; \ 0]$, and when $x \rightarrow -\infty, \phi = 1$ give $[1 \ -2 \ 3 \ -4 \ 5 \ ; \ 1]$.

After the augmented matrices of the system and conditions are computed, and the inverse of W^* determined, we obtain the coefficients vector as

$$A = \begin{bmatrix} \frac{21}{128} & \frac{-3}{16} & \frac{27}{256} & \frac{-1}{32} & \frac{1}{256} \end{bmatrix},$$

then, the solution is

$$\phi(x) = \frac{21}{128}E_0^U(x) - \frac{3}{16}E_1^U(x) + \frac{27}{256}E_2^U(x) - \frac{1}{32}E_3^U(x) + \frac{1}{256}E_4^U(x),$$

after simplifying the previous result we get

$$\phi(x) = \frac{1}{(1 + e^x)^4},$$

which is the exact solution of the problem (34), (see [28]).

Example 5.2. Consider the following differential equation [28,29,32]

$$\Im \phi = f(x), x \in (-\infty, \infty) \tag{35}$$

where, the operator $\Im = [\frac{d^2}{dx^2} - 1]$, and the subjected boundary conditions is $\phi(x) \rightarrow 0$ when $|x| \rightarrow \infty$.

The analytic exact solution given in [32] by Fourier transform as

$$\phi(x) = F^{-1} \left[\frac{-1}{1 + \omega^2} F[f(x)] \right],$$

where, F and F^{-1} are the Fourier and inverse Fourier transform operators, and the function $f(x)$ is well defined with respect to the Fourier transform conditions. We apply our present method to Eq. (35), by taking $f(x) = -2\text{sech}^3(x)$, the approximate and the exact solutions are compared as given in Table 1 at different N , where $x \in [-3, 3]$. The computing of the error norms L_2 and L_∞ given in Table 2 (by taking $h=0.1$), where

$$L_2 = \sqrt{h \sum_{i=0}^I (\phi_{Exact}^i - \phi_{Approximat}^i)^2},$$

$$L_\infty = \text{Max} |\phi_{Exact}^i - \phi_{Approximat}^i|$$

Fig. 1 show the approximate and exact solutions at different N , and $x \in [-10, 10]$, while Fig. 2 shows that the error function at different N and $x \in [0, 3]$

Example 5.3. Now, we consider the following problem [28,29]

$$\Re \phi = g(x), x \in (-\infty, \infty), \tag{36}$$

where, in this example the differential linear operator is $\Re = [\frac{d^3}{dx^3} - 1]$, (third order differential equation) and the boundary conditions is $\phi(x) \rightarrow \pm 1$ when $x \rightarrow \pm\infty$.

The exact solution taken as $\phi(x) = \tanh(x)$, and the function $g(x)$ takes the form

$$g(x) = -2\text{sech}^4(x) - \tanh(x) + 4\text{sech}^2(x)\tanh^2(x).$$




Fig. 1. The approximate and exact solutions at different N for Example 5.2.




Fig. 2. Error function of Example 5.2 at N=10, 12 and 16.

Table 3
The L_2 , L_∞ error norms for Example 5.3.

	L_2	L_∞
N = 10	1.15884×10^{-5}	1.42591×10^{-3}
N = 12	3.49288×10^{-9}	3.1855×10^{-5}
N = 16	8.68131×10^{-11}	5.24761×10^{-6}

by applying our present method to Eq. (36), the fundamental matrix is $\{Q_0V^{(0)}(M^T)^0 + Q_1V^{(1)}(M^T)^1 + Q_2V^{(2)}(M^T)^2 + Q_3V^{(3)}(M^T)^3\}A = G$,

Table 3 shows the L_2 and L_∞ error norms at different N where $h=0.1$ and $x \in [-3, 3]$, and Fig. 3 shows the approximate and exact solutions, and in Fig. 4 the error function at different N are given, $x \in [0, 3]$.

Example 5.4. Finally, we consider the fifth order differential equation as

$$\phi^{(5)} + \text{Sech}(x) \phi = \theta(x), \quad x \in (-\infty, \infty) \tag{37}$$

where, the conditions are in the following form

$$\begin{aligned} \phi(x) &= 1 \text{ when } x \rightarrow \infty \text{ and } \phi(x) = -1 \text{ when } x \rightarrow -\infty, \\ \phi(0) + 3\phi'(0) &= 3/2, \\ \phi''(0) &= 0. \end{aligned}$$

The fundamental matrix for the pervious equation is

$$\{Q_0V^{(0)}(M^T)^0 + Q_5V^{(5)}(M^T)^5\}A = \Theta,$$

where, $Q_1 = Q_2 = Q_3 = Q_4 = 0$, the exact solution taken as $\phi(x) = \tanh(x/2)$, and the function $\theta(x)$ will be in the form

$$\theta(x) = \frac{1}{16} \{ (33 - 26 \cosh(x) + \cosh(2x)) \text{sech}^6(x/2) + 16 \tanh(x/2) \text{sech}(x) \}.$$

The solution with the present method where $N=8$ obtained as

$$A = \begin{bmatrix} 0 & \frac{1}{2} & 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix},$$

then,

$$\phi(x) = \frac{1}{2} E_1^U(x) = \frac{1}{2} \left(2 \left(\frac{e^x - 1}{e^x + 1} \right) \right) = \left(\frac{e^x - 1}{e^x + 1} \right) = \tanh(x/2),$$

which is the same with the exact solution of the problem (37).

6. Application

Our application is evaluating Dawson's integral by solving its differential equation using our introduced method.

Dawson's integral is defined by

$$u(x) = e^{-x^2} \int_0^x e^{t^2} dt. \tag{38}$$

Dawson's integral can be written in terms of the error function of imaginary argument as

$$u(x) = -e^{-x^2} \frac{i}{2} \sqrt{\pi} \text{erf}(ix). \tag{39}$$

Dawson's integral is important and has many applications, there are many reports on Dawson's integral approximation [33–36].

In addition, the plasma dispersion function or "Faddeeva function" is Dawson's integral. Also, evaluating or approximating the error function in the complex plane is reported [37] which is an implicitly method for evaluating Dawson's integral. The differential equation corresponding to Dawson's integral is first order differential equation with boundary conditions tends to infinity, in the following form

$$u' + 2xu = 1, \quad x \in [-\infty, \infty]. \tag{40}$$

With the subjected conditions are that $u(x)$ is bounded (equal zero), as $|x| \rightarrow \infty$, Eq. (40) is non-homogenous first order boundary value problem. The square bracket in Eq. (40) which contains the infinity seems to be right because of x already tends to $\pm \infty$. Boyd [36] also used the previous differential equation to approximate Dawson's integral using rational function expansion in terms of Chebyshev polynomial of second kind.

Now we apply our proposed method to solve Eq. (40) with subjected conditions.

The fundamental matrix of Eq. (40) is

$$\{Q_0V^{(0)}(M^T)^0 + Q_1V^{(1)}(M^T)^1\}A = F,$$

After simplifying and finding the approximate solution as pervious examples with $N = 10, 16$, and 24 , the numerical results obtained as follows.

In Table 4 the L_2, L_∞ error norms show at the greater N gives lower error, the computation compared with the exact solution




Fig. 3. The approximate and exact solution at different N for Example 5.3.




Fig. 4. Error function of Example 5.3 at $N=10, 14$ and 16 .




Fig. 6. Error function of Dawson's integral at $N=10, 16$ and 24 .

Table 4
Comparing the L_2, L_∞ norms.

	L_2	L_∞
$N = 10$	5.88292×10^{-4}	1.06548×10^{-2}
$N = 16$	8.92815×10^{-5}	4.3758×10^{-3}
$N = 24$	1.60695×10^{-5}	2.1532×10^{-3}

given in (39) where $h=0.1$ and $x \in [-3, 3]$, in addition Fig. 5 shows the approximate and exact solutions where $N=16$ and 24 , also Fig. 6 shows the error function at different $N, x \in [0, 3]$

7. Conclusion

A new exponential Chebyshev of second kind (ESC) operational matrix of derivatives is investigated. The new operational matrix of

derivatives of the ESC functions is derived and introduced for solving high-order linear ordinary differential equations with variable coefficients in unbounded domains using the collocation method. The proposed differential equations and the given conditions are transformed to matrix equation with unknown ESC coefficients. On the other hand, the ESC functions approach deals directly with infinite boundaries without singularities or divergence. This variant for our method gave us freedom to solve differential equations with boundary conditions tend to infinity. Illustrative examples are used to demonstrate the applicability and the effectiveness of the proposed technique. As an application of our method approximating Dawson's integral by solving its differential equation is introduced. The corresponding differential equation to Dawson's integral is first order boundary value problem with conditions tends to infinity. The numerical results give good accuracy after comparing with the




Fig. 5. The approximate and exact solution at different N for Dawson's integral.

exact solution. The method can be extended for the case of non-linear ordinary, systems of linear differential and intgro-differential equations with variable coefficients which is under investigation by the authors as future work.

Acknowledgment

The authors sincerely thank the editors, reviewers and everyone who provide the advice, support, help and useful comments. The product of this research paper would not be possible without all of them.

Appendix

Code for Ex.5.1

```
n=4;
v[x_]=(e^x-1)/(e^x+1);
q[0,x_]=(-15*e^2*x)/(1+e^x)^2;
q[1,x_]=-1/(1+e^x);
q[2,x_]=1;
f[x_]=e^2x/(1+e^x)^6;
For[i=0,i ≤ n,i++,x[i]=Log[(1+Cos[(i*π)/n])/(1-Cos[(i*π)/n])]];
q0=Array[q00,{n+1,n+1},{0,0}];
For[i=0,i ≤ n,i++,For[j=0,j ≤ n,j++,q00[i,j]=0]]
For[i=0,i ≤ n,i++,q00[i,i]=Limit[q[0,x],x → x[i]]]
MatrixForm[q0]
q1=Array[q11,{n+1,n+1},{0,0}];
For[i=0,i ≤ n,i++,For[j=0,j ≤ n,j++,q11[i,j]=0]]
For[i=0,i ≤ n,i++,q11[i,i]=Limit[q[1,x],x → x[i]]]
MatrixForm[q1]
q2=Array[q22,{n+1,n+1},{0,0}];
For[i=0,i ≤ n,i++,For[j=0,j ≤ n,j++,q22[i,j]=0]]
For[i=0,i ≤ n,i++,q22[i,i]=q[2,x[i]]]
MatrixForm[q2]
v0=Array[v00,{n+1,n+1},{0,0}];
For[i=1,i ≤ n+1,i++,For[j=1,j ≤ n+1,j++,v00[i,j]=0]]
For[i=0,i ≤ n,i++,v00[i,0]=1]
For[i=1,i ≤ n,i++,For[j=0,j ≤ n,j++,v00[j,i]=Limit[(v[x])^i,x → x[j]]]
MatrixForm[v0]
v1=Array[v11,{n+1,n+1},{0,0}];
For[i=1,i ≤ n+1,i++,For[j=1,j ≤ n+1,j++,v11[i,j]=0]]
For[i=0,i ≤ n,i++,v11[i,0]=0]
For[i=1,i ≤ n,i++,For[j=0,j ≤ n,j++,v11[j,i]=Limit[D[(v[x])^i,x → x[j]]]
MatrixForm[v1]
v2=Array[v22,{n+1,n+1},{0,0}];
For[i=1,i ≤ n+1,i++,For[j=1,j ≤ n+1,j++,v22[i,j]=0]]
For[i=0,i ≤ n,i++,v22[i,0]=0]
For[i=1,i ≤ n,i++,For[j=0,j ≤ n,j++,v22[j,i]=Limit[D[D[(v[x])^i,x → x[j]]]
MatrixForm[v2]
m=Array[m1,{n+1,n+1},{0,0}];
For[i=0,i ≤ n+1,i++,For[j=0,j ≤ n+1,j++,m1[i,j]=0]]
m1[0,0]=1;
For[i=1,i ≤ n,i++,If[OddQ[i],For[j=0,j ≤ (i-1)/2,j++,m1[i,i-2*j]=(-1)^j*
2^{i-2*j}*Binomial[(i-j),j]];If[EvenQ[i],For[j=0,j ≤ i/2,j++,m1[i,i-2*j]=(-
1)^j* 2^{i-2*j}*Binomial[(i-j),j]]]
MatrixForm[m]
mt=Transpose[m]
{{1,0,-1,0,1},{0,2,0,-4,0},{0,0,4,0,-12},{0,0,0,8,0},{0,0,0,0,16}}
va=Array[va1,n+1,0];
For[i=0,i ≤ n,i++,va1[i]=Limit[(v[x])^i,x → ∞]]
va.mt
```

```
{1,2,3,4,5}
vb=Array[vb1,n+1,0];
For[i=0,i ≤ n,i++,vb1[i]=Limit[(v[x])^i,x → -∞]]
vb.mt
{1,-2,3,-4,5}
f1=Array[f11,n+1,0];
For[i=0,i ≤ n,i++,f11[i]=Limit[f[x],x → x[i]]]
q0.v0.mt+q1.v1.mt+q2.v2.mt
w={ { % },va.mt, vb.mt}
wInv=Inverse[w]
Simplify[wInv.f1]
{21/128,-(3/16),27/256,-(1/32),1/256}
Simplify[21/128 ChebyshevU[0,(e^x-1)/(e^x+1)]-3/16 ChebyshevU
[1,(e^x-1)/(e^x+1)]+27/256 ChebyshevU[2,(e^x-1)/(e^x+1)]-1/32
ChebyshevU[3,(e^x-1)/(e^x+1)]+1/256 ChebyshevU[4,(e^x-1)/(e^x
+1)]]
1/(1+e^x)^4
```

Code for Ex.5.4

```
n=8;
v[x_]=(e^x-1)/(e^x+1);
q[0,x_]=Sech[x];
q[5,x_]=1;
g[x_]=1/16 ((33-26 Cosh[x]+Cosh[2 x]) Sech[x/2]^6+16 Sech[x] Tanh
[x/2]);
For[i=0,i ≤ n,i++,x[i]=Log[(1+Cos[(i*π)/n])/(1-Cos[(i*π)/n])]];
q0=Array[q00,{n+1,n+1},{0,0}];
For[i=0,i ≤ n,i++,For[j=0,j ≤ n,j++,q00[i,j]=0]]
For[i=0,i ≤ n,i++,q00[i,i]=Limit[q[0,x],x → x[i]]]
MatrixForm[q0]
q5=Array[q55,{n+1,n+1},{0,0}];
For[i=0,i ≤ n,i++,For[j=0,j ≤ n,j++,q55[i,j]=0]]
For[i=0,i ≤ n,i++,q55[i,i]=Limit[q[5,x],x → x[i]]]
MatrixForm[q5]
v5=Array[v55,{n+1,n+1},{0,0}];
For[i=1,i ≤ n+1,i++,For[j=1,j ≤ n+1,j++,v55[i,j]=0]]
For[i=0,i ≤ n,i++,v55[i,0]=0]
For[i=1,i ≤ n,i++,For[j=0,j ≤ n,j++,v55[j,i]=Limit[D[D[D[D[(v[x])^i,
x],x],x],x],x → x[j]]]
MatrixForm[v5]
m=Array[m1,{n+1,n+1},{0,0}];
For[i=0,i ≤ n+1,i++,For[j=0,j ≤ n+1,j++,m1[i,j]=0]]
m1[0,0]=1;
For[i=1,i ≤ n,i++,If[OddQ[i],For[j=0,j ≤ (i-1)/2,j++,m1[i,i-2*j]=(-1)^j*
2^{i-2*j}*Binomial[(i-j),j]];If[EvenQ[i],For[j=0,j ≤ i/2,j++,m1[i,i-2*j]=(-
1)^j*2^{i-2*j}*Binomial[(i-j),j]]]
MatrixForm[m]
mt=Transpose[m]
va=Array[va1,n+1,0];
For[i=0,i ≤ n,i++,va1[i]=Limit[(v[x])^i,x → ∞]]
va.mt
{1,2,3,4,5,6,7,8,9}
vb=Array[vb1,n+1,0];
For[i=0,i ≤ n,i++,vb1[i]=Limit[(v[x])^i,x → -∞]]
vb.mt
{1,-2,3,-4,5,-6,7,-8,9}
vc=Array[vc1,n+1,0];
For[i=0,i ≤ n,i++,vc1[i]=Limit[(v[x])^i,x → 0]]
vc.mt
{1,0,-1,0,1,0,-1,0,1}
vd=Array[vd1,n+1,0];
For[i=0,i ≤ n,i++,vd1[i]=Limit[D[(v[x])^i,x → 0]]]
vd.mt
{0,1,0,-2,0,3,0,-4,0}
```



```

ve=Array[ve1,n+1,0];
For[i=0,i ≤ n,i++,ve1[i]=Limit[D[D[(v[x])i,x],x],x → 0]]
ve.mt
{0,0,2,0,-6,0,12,0,-20}
g1=Array[g11,n+1,0];
For[i=0,i ≤ n,i++,g11[i]=Limit[g[x],x → x[i]]]
q0.v0.mt+q5.v5.mt
w={%,va.mt, vb.mt,(vc.mt+3*vd.mt),ve.mt}
wInv=Inverse[w]
Simplify[wInv.g1]
{0,0.5,0,0,0,0,0,0}
FullSimplify[0.5* ChebyshevU[1,(ex-1)/(ex+1)]
Tanh[x/2]

```

References

- [1] I.H. Siyyam, Laguerre tau methods for solving higher-order ordinary differential equations, *J. Comput. Anal. Appl.* 3 (2) (2001) 173–182.
- [2] K. Parand, M. Razzaghi, Rational Chebyshev tau method for solving higher-order ordinary differential equations, *Int. J. Comput. Math.* 81 (1) (2004) 73–80.
- [3] B. Guo, J. Yan, Legendre–Gauss collocation method for initial value problems of second order ordinary differential equations, *Appl. Numer. Math.* 59 (6) (2009) 1386–1408.
- [4] Z. Wang, B. Guo, Legendre–Gauss–Radau collocation method for solving initial value problems of first order ordinary differential equations, *J. Sci. Comput.* 52 (1) (2012) 226–255.
- [5] D.O. Awoyemi, O.M. Idowu, A class of hybrid collocation methods for third-order ordinary differential equations, *Int. J. Comput. Math.* 82 (10) (2005) 1287–1293.
- [6] S. Adjerid, H. Temimi, A discontinuous Galerkin method for higher-order ordinary differential equations, *Comput. Methods Appl. Mech. Eng.* 197 (1) (2007) 202–218.
- [7] C.R. Smith, The Sinc-Galerkin method for fourth-order differential equations, *SIAM J. Numer. Anal.* 28 (3) (1991) 760–788.
- [8] W.M. Abd-Elhameed, Hany, M. Ahmed, Y.H. Youssri, A new generalized Jacobi Galerkin operational matrix of derivatives: two algorithms for solving fourth-order boundary value problems, *Adv. Difference Equ.* 1 (2016) (2016) 1–16.
- [9] W.M. Abd-Elhameed, E.H. Doha, Y.H. Youssri, Efficient spectral-Petrov-Galerkin methods for third-and fifth-order differential equations using general parameters generalized Jacobi polynomials, *Quaestiones Mathematicae* 36 (1) (2013) 15–38.
- [10] W.M. Abd-Elhameed, Y.H. Youssri, E.H. Doha, A novel operational matrix method based on shifted Legendre polynomials for solving second-order boundary value problems involving singular, singularly perturbed and Bratu-type equations, *Math. Sci.* 9 (2) (2015) 93–102.
- [11] E.H. Doha, W.M. Abd-Elhameed, A.H. Bhrawy, New spectral-Galerkin algorithms for direct solution of high even-order differential equations using symmetric generalized Jacobi polynomials, *Collectanea Mathematica* 64 (3) (2013) 373–394.
- [12] J.C. Mason, D.C. Handscomb, *Chebyshev Polynomials*, CRC Press, Boca Raton, 2003.
- [13] K. Wright, Chebyshev collocation methods for ordinary differential equations, *Comput. J.* 6 (4) (1964) 358–365.
- [14] M. Sezer, M. Kaynak, Chebyshev polynomial solutions of linear differential equations, *Int. J. Math. Edu. Sci. Technol.* 27 (4) (1996) 607–618.
- [15] J.P. Boyd, Orthogonal rational functions on a semi-infinite interval, *J. Comput. Phys.* 70 (1987) 63–88.
- [16] J.P. Boyd, Spectral methods using rational basis functions on an infinite interval, *J. Comput. Phys.* 69 (1987) 112–142.
- [17] J.P. Boyd, *Chebyshev and Fourier Spectral Methods*, Second ed., DOVER Publications, Mineola, 2000.
- [18] K. Parand, M. Razzaghi, Rational Chebyshev tau method for solving high-order ordinary differential equations, *Int. J. Comput. Math.* 81 (2004) 73–80.
- [19] K. Parand, A. Taghavi, M. Shahini, Comparison between rational Chebyshev and modified generalized Laguerre functions pseudospectral methods for solving Lane-Emden and unsteady gas equations, *Acta Phys. Pol. B* 40 (6) (2009) 1749.
- [20] M. Sezer, M. Gulsu, B. Tanay, Rational Chebyshev collocation method for solving higher-order linear ordinary differential equations, *Wiley Online Lib.* (2010), doi:10.1002/num.20573.
- [21] S. Yalcinbas, N. Ozsoy, M. Sezer, Approximate solution of higher-order linear differential equations by means of a new rational Chebyshev collocation method, *Math. Comput. Appl.* 15 (1) (2010) 45–56.
- [22] M.A. Ramadan, K.R. Raslan, M.A. Nassar, An approximate analytical solution of higher-order linear differential equations with variable coefficients using improved rational Chebyshev collocation method, *Appl. Comput. Math.* 3 (6) (2014) 315–322.
- [23] M.A. Ramadan, K.R. Raslan, M.A. Nassar, Numerical solution of system of higher order linear ordinary differential equations with variable coefficients using two proposed schemes for rational Chebyshev functions, *Global J. Math.* 3 (2) (2015) 322–327.
- [24] M.A. Ramadan, K.R. Raslan, M.A. Nassar, An approximate solution of systems of high-order linear differential equations with variable coefficients by means of a rational Chebyshev collocation method, *Electron. J. Math. Anal. Appl.* 4 (1) (2016) 86–98.
- [25] M.A. Ramadan, K.R. Raslan, A.R. Hadhoud, M.A. Nassar, Rational Chebyshev functions with new collocation points in semi-infinite domains for solving higher-order linear ordinary differential equations, *J. Adv. Math.* 7 (2015) 5403–5410.
- [26] M.A. Ramadan, K.R. Raslan, A.R. Hadhoud, M.A. Nassar, Numerical solution of high-order linear integro-differential equations with variable coefficients using two proposed schemes for rational Chebyshev functions, *New Trends in Mathematical Sciences* 4 (3) (2016) 22–35.
- [27] A.B. Koc, A. Kurnaz, A new kind of double Chebyshev polynomial approximation on unbounded domains, *Boundary Value Prob.* 1 (2013) (2013) 1687–2770.
- [28] M.A. Ramadan, K.R. Raslan, T.S. El Danaf, M.A. Abd El salam, On the exponential Chebyshev approximation in unbounded domains: a comparison study for solving high-order ordinary differential equations, *Int. J. Pure Appl. Math.* 105 (3) (2015) 399–413.
- [29] M.A. Ramadan, K.R. Raslan, T.S. El Danaf, M.A. Abd El salam, A new exponential Chebyshev operational matrix of derivatives for solving high-order ordinary differential equations in unbounded domains, *J. Mod. Methods Numer. Math.* 7 (1) (2016) 19–30.
- [30] J.C. Mason, Chebyshev polynomials of second, third and fourth kinds in approximation, indefinite integration, and integral transforms, *J. Comput. Appl. Math.* 49 (1993) 169–178.
- [31] N.H. Sweilam, A.M. Nagy, Adel, A. El-Sayed, Second kind shifted Chebyshev polynomials for solving space fractional order diffusion equation, *Chaos Solit. Fract.* 73 (2015) 141–147.
- [32] A.C. King, J. Billingham, S.R. Otto, *Differential Equations Linear, Nonlinear, Ordinary, Partial*, Cambridge University Press, 2003.
- [33] W.J. Cody, K.A. Paciorek, H.C. Thacher, Chebyshev approximations for Dawson's integral, *Math. Comput.* 23 (1970) 171–178.
- [34] J.P. Coleman, Complex polynomial approximation by the Lanczos-s-method: Dawson's integral, *J. Comput. Appl. Math.* 20 (1987) 137–151.
- [35] F.G. Lether, Constrained near-minimax rational approximations to Dawson's integral, *Appl. Math. Comput.* 88 (1997) 267–274.
- [36] J.P. Boyd, Evaluating of Dawson's Integral by solving its differential equation using orthogonal rational Chebyshev functions, *Appl. Math. Comput.* 204 (2008) 914–919.
- [37] J.A.C. Weideman, Computation of the complex error function, *SIAM J. Numer. Anal.* 31 (1994) 1497–1518.