



Egyptian Mathematical Society  
**Journal of the Egyptian Mathematical Society**

www.etms-eg.org  
 www.elsevier.com/locate/joems



ORIGINAL ARTICLE

# On a discretization process of fractional-order Logistic differential equation



Z.F. El Raheem, S.M. Salman \*

Faculty of Education, Alexandria University, Alexandria, Egypt

Received 12 July 2013; revised 12 September 2013; accepted 24 September 2013

Available online 14 November 2013

**KEYWORDS**

Logistic differential equation;  
 Piecewise constant arguments;  
 Fractional-order differential equations;  
 Fixed points;  
 Lyapunov exponent;  
 Bifurcation

**Abstract** In this work we introduce a discretization process to discretize fractional-order differential equations. First of all, we consider the fractional-order Logistic differential equation then, we consider the corresponding fractional-order Logistic differential equation with piecewise constant arguments and we apply the proposed discretization on it. The stability of the fixed points of the resultant dynamical system and the Lyapunov exponent are investigated. Finally, we study some dynamic behavior of the resultant systems such as bifurcation and chaos.

**2010 MATHEMATICS SUBJECT CLASSIFICATION:** 39B05; 37N30; 37N20

© 2013 Production and hosting by Elsevier B.V. on behalf of Egyptian Mathematical Society.  
 Open access under [CC BY-NC-ND license](#).

**1. Introduction**

Chaotic systems have been a focal point of renewed interest for many researchers in the past few decades. Such nonlinear systems can occur in various natural and man-made systems, and are known to have great sensitivity to initial conditions. In recent years differential equations with fractional-order have attracted many researchers because of their applications in many areas of science and engineering. Analytical and numerical techniques have been implemented to study such equations. The fractional calculus has allowed the operations of integra-

tion and differentiation to be applied upon any fractional-order. For the existence of solutions for fractional differential equations, one can see [1,2].

About the development of existence theorems for fractional functional differential equations, many contributions existed and can be referred to [3–5]. Many applications of fractional calculus amounts to replace the time derivative in a given evolution equation by a derivative of fractional-order.

Recalling the basic definitions (Caputo) and properties of fractional-order differentiation and integration

**Definition 1.** The fractional integral of order  $\beta \in \mathbb{R}^+$  of the function  $f(t)$ ,  $t > 0$  is defined by

$$I^\beta f(t) = \int_0^t \frac{(t-s)^{\beta-1}}{\Gamma(\beta)} f(s) ds,$$

and the fractional derivative of order  $\alpha \in (n-1, n)$  of  $f(t)$ ,  $t > 0$  is defined by

\* Corresponding author. Tel.: +20 1091103980.

E-mail addresses: [zaki55@hotmail.com](mailto:zaki55@hotmail.com) (Z.F. El Raheem), [samastars9@gmail.com](mailto:samastars9@gmail.com) (S.M. Salman).

Peer review under responsibility of Egyptian Mathematical Society.



Production and hosting by Elsevier

$$D^\alpha f(t) = I^{n-\alpha} D^n f(t), \quad D = \frac{d}{dt}.$$

To solve fractional-order differential equations there are two famous methods: frequency domain methods [6] and time domain methods [7]. In recent years it has been shown that the second method is more effective because the first method is not always reliable in detecting chaos [8,9].

Often it is not desirable to solve a differential equation analytically, and one turns to numerical or computational methods.

In [10], a numerical method for nonlinear fractional-order differential equations with constant or time-varying delay was devised. It should be noticed that the fractional differential equations tend to lower the dimensionality of the differential equations in question, however, introducing delay in differential equations makes it infinite dimensional. So, even a single ordinary differential equation with delay could display chaos.

On the other hand, some examples of dynamical systems generated by piecewise constant arguments have been studied in [11–14]. Here we propose a discretization process to obtain the discrete version of the system under study. Mean while, we apply discretization process to discretize the fractional-order Logistic differential equation.

A lot of differential equations with Caputo fractional derivative were simulated by the Predictor-Corrector scheme, such as the fractional Chua system, the fractional Chen system, and Lorenz system. We should note that Predictor-Corrector method is an approximation for the fractional-order integration, however, our approach is an approximation for the right hand side. For applications of fractional-order differential equations one can see [15–17], [21], and [23–27].

### 2. Discretization process

Consider the fractional-order Logistic differential equation given by

$$D^\alpha x(t) = \rho x(t)(1 - x(t)), \quad t > 0, \tag{2.1}$$

with the initial condition  $x(0) = x_0$ .

The main purpose of this section is to introduce a discretization process to discretize the counterpart of (2.1) with piecewise constant arguments

$$D^\alpha x(t) = \rho x\left(\left[\frac{t}{r}\right]r\right)\left(1 - x\left(\left[\frac{t}{r}\right]r\right)\right), \tag{2.2}$$

with the initial condition  $x(0) = x_0$ .

We proceed like the step method mentioned in [20] and [22]. The steps of the discretization process is as follows

(1) Let  $t \in [0, r)$ , then  $\frac{t}{r} \in [0, 1)$ . So, we get

$$D^\alpha x(t) = \rho x_0(1 - x_0), \quad t \in [0, r),$$

and the solution of (2.2) is given by

$$\begin{aligned} x_1(t) &= x_0 + I^\alpha \rho x_0(1 - x_0) = x_0 + \rho x_0(1 - x_0) \int_0^t \frac{(t-s)^{\alpha-1}}{\Gamma(\alpha)} ds \\ &= x_0 + \rho x_0(1 - x_0) \frac{t^\alpha}{\Gamma(1 + \alpha)} \end{aligned}$$

(2) Let  $t \in [r, 2r)$ , then  $\frac{t}{r} \in [1, 2)$ . So, we get

$$D^\alpha x(t) = \rho x_1(1 - x_1), \quad t \in [r, 2r),$$

and the solution of (2.2) is given by

$$\begin{aligned} x_2(t) &= x_1(r) + I_r^\alpha \rho x_1(1 - x_1) \\ &= x_1(r) + \rho x_1(1 - x_1) \int_r^t \frac{(t-s)^{\alpha-1}}{\Gamma(\alpha)} ds \\ &= x_1(r) + \rho x_1(r)(1 - x_1(r)) \frac{(t-r)^\alpha}{\Gamma(1 + \alpha)} \end{aligned}$$

Repeating the process we can easily deduce that the solution of (2.2) is given by

$$\begin{aligned} x_{n+1}(t) &= x_n(nr) + \frac{(t - nr)^\alpha}{\Gamma(1 + \alpha)} \rho x_n(nr)(1 - x_n(nr)), \quad t \\ &\in [nr, (n + 1)r). \end{aligned}$$

Let  $t \rightarrow (n + 1)r$ , we obtain the discretization

$$x_{n+1}((n + 1)r) = x_n(nr) + \frac{r^\alpha}{\Gamma(1 + \alpha)} \rho x_n(nr)(1 - x_n(nr)),$$

That is

$$x_{n+1} = x_n + \frac{r^\alpha}{\Gamma(1 + \alpha)} \rho x_n(1 - x_n). \tag{2.3}$$

On a similar manner, consider the corresponding equation of (2.1) with piecewise constant arguments

$$D^\alpha x(t) = \rho x\left(\left[\frac{t}{r}\right]r\right)\left(1 - x\left(\left[\frac{t}{r}\right]r\right)\right), \tag{2.4}$$

with the initial condition  $x(0) = x_0$ . So, we obtain the second order discretization

$$x_{n+1} = x_n + \frac{r^\alpha}{\Gamma(1 + \alpha)} \rho x_n(1 - x_{n-1}). \tag{2.5}$$

### 3. Fixed points and their asymptotic stability

Now we study the stability of the fixed points of the Eq. (2.3) which has two fixed points namely, 0 and 1 given by solving the equation

$$x = x + \frac{r^\alpha}{\Gamma(1 + \alpha)} \rho x(1 - x).$$

To study the stability of these fixed points we rely on the following theorem

**Theorem 1** [18]. *Let  $f$  be a smooth map on  $\mathbb{R}$ , and assume that  $x_0$  is a fixed point of  $f$ .*

1. If  $-f'(x_0) < 1$ , then  $x_0$  is stable.
2. If  $-f'(x_0) > 1$ , then  $x_0$  is unstable.

In case of the first fixed point ‘0’, it is stable if  $\left|1 + \frac{r^\alpha}{\Gamma(1 + \alpha)} \rho\right| < 1$  which is impossible. That is the origin is unstable. For the second fixed point ‘1’, it is stable if

$$0 < \rho < \frac{2(\Gamma(1 + \alpha))}{r^\alpha}. \tag{3.1}$$

On the other hand, to study the stability of the fixed points of Eq. (2.5) we first split it into two equations as follows






Fig. 1 Lyapunov exponent for system (2.3) with different values of the fractional-order parameter  $\alpha$ .



**Fig. 2** Bifurcation diagram of system (3.2) as a function of  $\rho$  with different values of the fractional-order parameter  $\alpha$ .



**Fig. 3** Bifurcation diagram of system (3.2) as a function of the fractional-order parameter  $\alpha$  with different values of the parameters  $r$  and  $\rho$ .

$$\begin{aligned}
 y_{n+1} &= x_n \\
 x_{n+1} &= x_n + \frac{r^\alpha}{\Gamma(1+\alpha)} \rho x_n (1 - y_n).
 \end{aligned}
 \tag{3.2}$$

This system has two fixed points namely  $(x, y)_{fix1} = (0, 0)$  and  $(x, y)_{fix2} = (1, 1)$

By considering a Jacobian matrix for one of these fixed points and calculating their eigenvalues, we can investigate the stability of each fixed point based on the roots of the system characteristic equation [19]. The Jacobian matrix is given by

$$J = \begin{pmatrix} 1 & 0 \\ 1 + \frac{r^\alpha \rho}{\Gamma(1+\alpha)} (1 - y) & -\frac{r^\alpha \rho}{\Gamma(1+\alpha)} x \end{pmatrix}$$

The eigenvalues associated to the Jacobian matrix for the first fixed point are  $\lambda_1 = 0$ , and  $\lambda_2 = 1$ , that is, this fixed point is unstable.

While the eigenvalues associated to the Jacobian matrix for the second fixed point are

$$\begin{aligned}
 \lambda_{1,2} &= 0.5 \left( 1 - \frac{r^\alpha \rho}{\Gamma(1+\alpha)} \right) \pm 0.5 \\
 &\quad \times \sqrt{\left( -1 + \frac{r^\alpha \rho}{\Gamma(1+\alpha)} \right)^2 + 4 \frac{r^\alpha \rho}{\Gamma(1+\alpha)}}.
 \end{aligned}$$

If we take for instance  $r = 0.2$ ,  $\alpha = 0.85$ , and  $\rho = 3$ , we get  $\lambda_1 = 1$  and  $\lambda_2 = -0.0539$ . This means that this fixed point is unstable.

In the next section, the numerical experiments assure our analytical results for different values for  $r$ ,  $\alpha$ , and  $\rho$ . It is worth to mention here that Lyapunov exponent for (2.3) is given by (see 1).

$$Lya.exp = \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \log_2 \left( 1 + \frac{r^\alpha}{\Gamma(1+\alpha)} \rho (1 - 2x) \right).$$

When  $\alpha \rightarrow 1$  the same Lyapunov exponent for the original discrete system ( $x_{n+1} = x_n + r \rho x_n (1 - x_n)$ ) is obtained. The following figures show the lyapunov exponent for the system (2.3) for different values of the fractional-order parameter  $\alpha$



#### 4. Bifurcation and chaos

In this section we show by numerical experiments bifurcation and chaos of the dynamical system (3.2) first with respect to the parameter  $\rho$  and then with respect to the fractional-order parameter  $\alpha$ .

Let  $r = 0.25$  be fixed and vary  $\alpha$  from 0.70 to 0.95 and  $\rho$  from 0 to 8. The initial state of the system (3.2) is  $x_0 = 0.1$  and  $y_0 = 0.2$ . The step size for  $\rho$  is 0.001, the resulting bifurcation diagrams are shown in Fig. (2) from (a) – (f). It is observed from the figures that increasing the fractional-order parameter  $\alpha$  and fixing the parameter  $r$  stabilize the chaotic system.

Now vary the fractional-order parameter  $\alpha$  from 0.70 to 0.95 but with a fixed system parameter  $\rho$  and change the parameter  $r$  from 0.15 to 0.30. The resulting bifurcation diagrams are shown in Fig. (3) from (a) – (d).

#### 5. Conclusion

In this work we studied the dynamics of the fractional-order Logistic equation. We applied a simple discretization scheme to discretize fractional-order differential equations. Chaos and bifurcation of the resulting discrete system were numerically investigated by varying the system parameter  $\rho$  and the fractional-order parameter  $\alpha$ . We have noticed that when  $\alpha \rightarrow 1$ , the discretization will be Euler's method discretization [18]. Moreover, Euler's method is able to discretize a first order difference equations, however, we succeeded in discretizing a second order difference equation.

#### Acknowledgement

The authors thank Prof. Dr. El-Sayed Ahmed and Prof. Dr. Ahmed Matouk for their discussion and valuable comments.

#### References

- [1] A. El-Sayed, A. El-Mesiry, H. EL-Saka, On the fractional-order logistic equation, *Applied Mathematics Letters* 20 (2007) 817–823.
- [2] A.M.A. El-Sayed, Nonlinear functional–differential equations of arbitrary orders, *Nonlinear Analysis* 33 (1998) 2 (181186).
- [3] S. Das, *Functional Fractional Calculus for System Identification and Controls*, Springer, 2007.
- [4] I. Podlubny, *Fractional Differential Equations*, Mathematics in Science and Engineering, vol. 198, Academic Press, San Diego, CA, USA, 1999.
- [5] I. Podlubny, Geometric and physical interpretation of fractional integration and fractional differentiation. Dedicated to the 60th anniversary of Prof. Francesco Mainardi, *Fraction Calculus in Application Analysis* 5 (4) (2002) 367–386.
- [6] H. Sun, A. Abdelwahed, B. Onaral, Linear approximation for transfer function with a pole of fractional-order, *IEEE Transactions on Automatic Control* 29 (1984) 441–444.
- [7] K. Diethelm, N.J. Ford, A.D. Freed, A predictor–corrector approach for the numerical solution of fractional differential equations, *Journal of Nonlinear Dynamics* 29 (2002) 3–22.
- [8] M.S. Tavazoei, M. Haeri, Unreliability of frequency domain approximation in recognizing chaos in fractional-order systems, *IET Signalling Process* 1 (2007) 171–181.
- [9] M.S. Tavazoei, M. Haeri, Limitation of frequency domain approximation for detecting chaos in fractional-order systems, *Journal of Nonlinear Analysis, Theory, Methods, & Applications* 69 (2008) 1299–1320.
- [10] Z. Wang, A numerical method for delayed fractional-order differential equations, *Journal of Applied Mathematics* 2013, 256071 (Hindawi, 7 pages).
- [11] M.U. Akhmet, Stability of differential equations with piecewise constant arguments of generalized type, *Nonlinear Analysis* 68 (2008) 4 794–803.
- [12] M.U. Akhmet, D. Altntana, T. Ergenc, Chaos of the logistic equation with piecewise constant arguments, arXiv:1006.4753, 2010.
- [13] A.M.A. El-Sayed, S.M. Salman, Chaos and bifurcation of discontinuous dynamical systems with piecewise constant arguments, *Malaya Journal of Matematik* 1 (1) (2012) 14–18.
- [14] A.M.A. El-Sayed, S.M. Salman, Chaos and bifurcation of the Logistic discontinuous dynamical systems with piecewise constant arguments, *Malaya Journal of Matematik* 3 (1) (2013) 14–20.
- [15] W. Guo-Cheng, B. Dumitru, New applications of the variational iteration method – from differential equations to q-fractional difference equations, *Advances in Difference Equations* 21 (0) (2013), doi:<http://dx.doi.org/10.1186/1687-1847-2013-21>.
- [16] V.D. Gejji, S. Bhalekar, P. Gade, Dynamics of fractional-order Chen systems with delays, *Pramana Journal of Physics* 79 (1) (2012) 61–69.
- [17] G.A. Khalili, Y.A. Moslemi, B. Dumitru, On the fractional Hamilton and Lagrange mechanics, *International Journal of Theoretical Physics* 51(9) (2012) 2909–2916, doi:<http://dx.doi.org/10.1007/s10773-012-1169-8>.
- [18] S.N. Elaidy, *An Introduction to Difference Equations, Third Edition., Undergraduate Texts in Mathematics*, Springer, New York, 2005.
- [19] R. Holmgren, *A First Course in Discrete Dynamical Systems*, Springer Verlag, New York, 1994.
- [20] N.A. Bohai, Continuous solutions of systems of nonlinear difference equations with continuous arguments and their properties, *Journal of Nonlinear oscillations* 10 (2) (2007).
- [21] Y.Q. Chen, B.M. Vinagre, I. Podlubny, A new discretization method for fractional-order differentiators via continued fraction expansion, in: *Proceedings of DETC*, vol. 3, pp. 761–769.
- [22] A.M.A. El-Sayed, S.M. Salman, On a discretization process of fractional-order Riccati differential equation, *Journal of Fractional Calculus and Applications* 4 (2) (2013) 251–259.
- [23] J. Fahd, A. Thabet, B. Dumitru, Stability of q-fractional non-autonomous systems, *Nonlinear Analysis – Real World Applications* 14 (1) (2013) 780–784, doi:<http://dx.doi.org/10.1016/j.nonrwa.2012.08.001>.
- [24] I. Podlubny, "Numerical solution of ordinary fractional differential equations by the fractional difference method", in: S. Elaydi, I. Gyori and G. Ladas, (Eds.), *Advances in Difference Equations*, Gordon and Breach, Amsterdam, 1997, pp. 507–516.
- [25] I. Podlubny, Matrix approach to discrete fractional calculus, *Fractional Calculus and Applied Analysis* 3 (4) (2000) 359–386.
- [26] I. Podlubny, The Laplace transform method for linear differential equations of the fractional-order, arXiv preprint [arXiv preprint funct-an/9710005](http://arxiv.org/abs/09710005).
- [27] B. Sachin, D. Varsha, B. Dumitru, et al, Generalized fractional-order Bloch equation with extended delay, *International Journal of Bifurcational and Chaos* 22 (4) (2012) 1–15, 1250071.