



Original Article

# Totally real submanifolds of Kaehler product manifolds



Majid Ali Choudhary

Department of Mathematics, Jamia Millia Islamia, New Delhi 110025, India

Received 22 September 2014; revised 28 December 2014; accepted 21 February 2015  
 Available online 23 May 2015

**Keywords**

Kaehlerian product manifold;  
 Totally real submanifold;  
 Complex space form

**Abstract** Totally real submanifolds have been studied by many geometers in different ambient manifolds. The purpose of this note is to study totally real submanifolds in Kaehlerian product manifolds. We derive some integral formulas computing the Laplacian of the square of the second fundamental form and using these formulas we prove pinching theorems. In fact, we have generalized some results due to Yano and Kon [1,2] to the case when the ambient manifold is Kaehlerian product manifold.

**2000 Mathematics Subject Classification:** 53C15; 53C40

Copyright 2015, Egyptian Mathematical Society. Production and hosting by Elsevier B.V.  
 This is an open access article under the CC BY-NC-ND license  
 (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

**1. Introduction**

The geometry of totally real submanifolds is an interesting field which is studied by many geometers. For example, Houh [3], Yau [4], Chen and Ogiue [5] have studied totally real submanifolds in an almost Hermitian manifold or a Kaehlerian manifold of constant holomorphic sectional curvature and obtained many interesting results. Moreover, Yano and Kon [1,2] have generalized some of the results proved in [6–9]. On the other hand, Kaehlerian product manifold has also been paid attention by geometers [10]. The object of this note is to study the

geometry of totally real submanifolds when the ambient manifold is a Kaehlerian product manifold.

**2. Preliminaries**

Let  $\bar{M}^n$  be a Kaehlerian manifold of complex dimension  $n$  (of real dimension  $2n$ ) and  $\bar{M}^p$  be a Kaehlerian manifold of complex dimension  $p$  (of real dimension  $2p$ ). Let us denote by  $J_n$  and  $J_p$  almost complex structures of  $\bar{M}^n$  and  $\bar{M}^p$  respectively. Now, we suppose that  $\bar{M}^n$  and  $\bar{M}^p$  are complex space forms with constant holomorphic sectional curvatures  $c_1$  and  $c_2$  and denote them by  $\bar{M}^n(c_1)$  and  $\bar{M}^p(c_2)$  respectively. The Riemannian curvature tensor  $\bar{R}_n$  of  $\bar{M}^n(c_1)$  is given by

$$\bar{R}_n(X, Y)Z = \frac{1}{4}c_1[g_n(Y, Z)X - g_n(X, Z)Y] + \frac{1}{4}c_1[g_n(J_n Y, Z)J_n X - g_n(J_n X, Z)J_n Y + 2g_n(X, J_n Y)J_n Z]$$

E-mail address: [majid\\_alichoudhary@yahoo.co.in](mailto:majid_alichoudhary@yahoo.co.in)  
 Peer review under responsibility of Egyptian Mathematical Society.



and the Riemannian curvature tensor  $\bar{R}_p$  of  $\bar{M}^p(c_2)$  is given by

$$\begin{aligned}\bar{R}_p(X, Y)Z &= \frac{1}{4}c_2[g_p(Y, Z)X - g_p(X, Z)Y] \\ &+ \frac{1}{4}c_2[g_p(J_p Y, Z)J_p X - g_p(J_p X, Z)J_p Y + 2g_p(X, J_p Y)J_p Z].\end{aligned}$$

We consider the Kaehlerian product manifold  $\bar{M} = \bar{M}^n(c_1) \times \bar{M}^p(c_2)$ . Let us denote by  $P$  and  $Q$  the projection operators of the tangent space of  $\bar{M}^n(c_1)$  and  $\bar{M}^p(c_2)$  respectively. Then, we have  $P^2 = P, Q^2 = Q, PQ = QP = 0$ . We put  $F = P - Q$  and it can be verified that  $F^2 = I$ . Thus,  $F$  is almost product structure on  $\bar{M}$ . Moreover, we define a Riemannian metric  $g$  on  $\bar{M}$  by

$$g(X, Y) = g_n(PX, PY) + g_p(QX, QY)$$

for any vector field  $X$  and  $Y$  of  $\bar{M}$ . It also follows that  $g(FX, Y) = g(FY, X)$ . Let us put  $JX = J_n PX + J_p QX$  for any vector field  $X$  of  $\bar{M}$ . Then we see that  $J_n P = PJ, J_p Q = QJ, FJ = JF, J^2 = -I, g(JX, JY) = g(X, Y), \bar{\nabla}_X J = 0$ . Thus,  $J$  is Kaehlerian structure on  $\bar{M}$ . The Riemannian curvature tensor  $\bar{R}$  of a Kaehlerian product manifold  $\bar{M}$  is given by [10]

$$\begin{aligned}\bar{R}(X, Y, Z, W) &= \frac{1}{16}(c_1 + c_2)[g(Y, Z)g(X, W) - g(X, Z)g(Y, W)] \\ &+ g(JY, Z)g(JX, W) - g(JX, Z)g(JY, W) \\ &+ 2g(X, JY)g(JZ, W) + 2g(FY, Z)g(FX, W) \\ &- g(FX, Z)g(FY, W) + g(FJY, Z)g(FJX, W) \\ &- g(FJX, Z)g(FJY, W) + 2g(FX, JY)g(FJZ, W)] \\ &+ \frac{1}{16}(c_1 - c_2)[g(FY, Z)g(X, W) - g(FX, Z)g(Y, W)] \\ &+ g(Y, Z)g(FX, W) - g(X, Z)g(FY, W) \\ &+ g(FJY, Z)g(JX, W) - g(FJX, Z)g(JY, W) \\ &+ g(JY, Z)g(FJX, W) - g(JX, Z)g(FJY, W) \\ &+ 2g(FX, JY)g(JZ, W) \\ &+ 2g(X, JY)g(JFZ, W)]\end{aligned}\quad (2.1)$$

for any vector fields  $X, Y$  and  $Z$  on  $\bar{M}$ . An  $n$ -dimensional Riemannian manifold  $M$  isometrically immersed in a Kaehlerian product manifold  $\bar{M}$  is called totally real submanifold of  $\bar{M}$  if  $JT_x(M) \perp T_x(M)$  for each  $x \in M$  where  $T_x(M)$  denotes the tangent space to  $M$  at  $x \in M$ . Here we have identified  $T_x(M)$  with its image under the differential of the immersion because our computation is local. If  $X \in T_x(M)$ , then  $JX$  is a normal vector to  $M$ . Let  $\bar{g}$  be the metric tensor field of  $\bar{M}$  and  $g$  be the induced metric tensor field on  $M$ . We denote by  $\bar{\nabla}$  (resp.  $\nabla$ ) the operator of covariant differentiation with respect to  $\bar{g}$  (resp.  $g$ ). Then the Gauss and Weingarten formulas are given by

$$\bar{\nabla}_X Y = \nabla_X Y + B(X, Y) \quad (2.2)$$

$$\bar{\nabla}_X N = -A_N X + D_X N \quad (2.3)$$

for any tangent vector fields  $X, Y$  and normal vector field  $N$  on  $M$ , where  $D$  is the operator of covariant differentiation with respect to the linear connection induced in the normal bundle. Both  $A$  and  $B$  are called the second fundamental form of  $M$  and satisfy

$$\bar{g}(B(X, Y), N) = g(A_N X, Y). \quad (2.4)$$

A normal vector field  $N$  in the normal bundle is said to be parallel if  $D_X N = 0$  for any tangent vector field  $X$  on  $M$ . The mean curvature vector  $H$  is defined as  $H = (1/n)TrB$ , where  $TrB = \sum_i B(e_i, e_i)$  for an orthonormal frame  $\{e_i\}$ . We say that

- $M$  is minimal if  $H = 0$ .
- $M$  is totally umbilical if the second fundamental form of  $M$  satisfies  $B(X, Y) = g(X, Y)H$ .
- $M$  is totally geodesic if the second fundamental form of  $M$  vanishes identically, that is,  $B = 0$ .

We choose a local field of orthonormal frames  $e_1, \dots, e_n; e_{n+1}, \dots, e_{n+p}; e_{1^*} = Je_1, \dots, e_{n^*} = Je_n; e_{(n+1)^*} = Je_{n+1}, \dots, e_{(n+p)^*} = Je_{n+p}$  in  $\bar{M}$  in such a way that restricted to  $M, e_1, \dots, e_n$  are tangent to  $M$ . With respect to this frame field of  $\bar{M}$ , let  $\omega^1, \dots, \omega^n; \omega^{n+1}, \dots, \omega^{n+p}; \omega^{1^*}, \dots, \omega^{n^*}; \omega^{(n+1)^*}, \dots, \omega^{(n+p)^*}$  be the field of dual frames. Unless otherwise stated, we use the conventions that the ranges of indices are respectively  $A, B, C, D = 1, \dots, n + p, 1^*, \dots, (n + p)^*; i, j, k, l, t, s = 1, \dots, n; a, b, c, d = n + 1, \dots, n + p, 1^*, \dots, (n + p)^*; \alpha, \beta, \gamma = n + 1, \dots, n + p; \lambda, \mu, \nu = n + 1, \dots, n + p, (n + 1)^*, \dots, (n + p)^*$  and that when an index appears twice in any term as a subscript and a superscript, it is understood that this index is summed over its range. Then the structure equations of  $\bar{M}$  are given by

$$\begin{aligned}d\omega^A &= -\omega_B^A \omega^B, \quad \omega_B^A + \omega_A^B = 0, \\ \omega_j^i + \omega_i^j &= 0, \quad \omega_j^i = \omega_{j^*}^{i^*}, \quad \omega_j^{i^*} = \omega_i^{j^*}, \\ \omega_\beta^\alpha + \omega_\alpha^\beta &= 0, \quad \omega_\beta^\alpha = \omega_{\beta^*}^{\alpha^*}, \quad \omega_\beta^{\alpha^*} = \omega_\alpha^{\beta^*},\end{aligned}\quad (2.5)$$

$$\begin{aligned}\omega_\alpha^i + \omega_i^\alpha &= 0, \quad \omega_\alpha^i = \omega_{\alpha^*}^{i^*}, \quad \omega_\alpha^{i^*} = \omega_i^{\alpha^*}, \\ d\omega_B^A &= -\omega_C^A \omega_B^C + \phi_B^A, \quad \phi_B^A = \frac{1}{2}K_{BCD}^A \omega^C \wedge \omega^D\end{aligned}\quad (2.6)$$

Restricting these forms to  $M$ , we have

$$\omega^a = 0, \quad (2.7)$$

$$d\omega^i = -\omega_k^i \wedge \omega^k, \quad (2.8)$$

$$d\omega_j^i = -\omega_k^i \wedge \omega_j^k + \Omega_j^i, \quad \Omega_j^i = \frac{1}{2}R_{jkl}^i \omega^k \wedge \omega^l \quad (2.9)$$

Since  $0 = d\omega^a = -\omega_i^a \wedge \omega^i$ , by Cartan's lemma we have

$$\omega_i^a = h_{ij}^a \omega^j, \quad h_{ij}^a = h_{ji}^a \quad (2.10)$$

We see that  $g(A_a e_i, e_j) = h_{ij}^a$ . The Gauss-equation is given by

$$R_{jkl}^i = K_{jkl}^i + \sum_a (h_{ik}^a h_{jl}^a - h_{il}^a h_{jk}^a) \quad (2.11)$$

Moreover we have

$$d\omega_b^a = -\omega_c^a \wedge \omega_b^c + \Omega_b^a, \quad \Omega_b^a = \frac{1}{2}R_{bkl}^a \omega^k \wedge \omega^l \quad (2.12)$$

and the Ricci-equation is given by

$$R_{bkl}^a = K_{bkl}^a + \sum_i (h_{ik}^a h_{il}^b - h_{il}^a h_{ik}^b) \quad (2.13)$$

From (2.5) and (2.10) we have

$$h_{jk}^{i^*} = h_{ik}^{j^*} = h_{ij}^{k^*} \quad (2.14)$$

We define the covariant derivative  $h_{ijk}^a$  of  $h_{ij}^a$  by setting

$$h^a_{ijk} \omega^k = dh^a_{ij} - h^a_{il} \omega^l_j - h^a_{lj} \omega^l_i + h^b_{ij} \omega^a_b. \tag{2.15}$$

The Laplacian  $\Delta h^a_{ij}$  of  $h^a_{ij}$  is defined as

$$\Delta h^a_{ij} = \sum_k h^a_{ijkk} \tag{2.16}$$

where we have put  $h^a_{ijkl} \omega^l = dh^a_{ijk} - h^a_{ljk} \omega^l_i - h^a_{ilk} \omega^l_j - h^a_{ijl} \omega^l_k + h^b_{ijk} \omega^a_b$ . The forms  $(\omega^a_j)$  define the Riemannian connection of  $M$  and the forms  $(\omega^a_b)$ . If a Riemannian manifold  $M$  is of constant curvaturbare  $k$ , then we have

$$R^i_{jkl} = k(\delta_{ik} \delta_{jl} - \delta_{il} \delta_{jk}). \tag{2.17}$$

We call such a manifold a real space form and denote it by  $M(k)$ .

Now, let us suppose that  $M$  be a totally real  $n$ -dimensional submanifold of a Kaehlerian product manifold  $\overline{M}$ . We denote by  $T_x(M)$  the tangent space of  $M$  at  $x \in M$  and by  $T_x(M)^\perp$  the normal space of  $M$  at  $x \in M$ . Then we see that  $JT_x(M) \subset T_x(M)^\perp$ . We can decompose  $T_x(M)^\perp$  in the following way:

$$T_x(M)^\perp = JT_x(M) \oplus N_x(M)$$

where  $N_x(M)$  is an orthogonal complement of  $JT_x(M)$  in  $T_x(M)^\perp$ . If  $N \in N_x(M)$ , we obtain  $JN \in N_x(M)$ . If  $N$  is a vector field in the normal bundle  $T(M)^\perp$ , we put

$$JN = PN + fN \tag{2.18}$$

where  $PN$  and  $fN$  are the tangential and normal part of  $JN$ . Then  $P$  is a tangent bundle valued 1-form on the normal bundle and  $f$  is an endomorphism of the normal bundle. Putting  $N = JX$  and applying  $J$  in (2.18), we find [7,9]:

$$PfN = 0, \quad f^2N = -N - JPN, \quad PJX = -X, \quad fJX = 0 \tag{2.19}$$

where  $X$  is a tangent vector field to  $M$  and  $N$  is a vector field in the normal bundle. Eq. (2.19) imply that

$$f^3 + f = 0.$$

Therefore,  $f$  being of constant rank, if  $f$  does not vanish, then it defines an  $f$ -structure in the normal bundle [11]. From (2.18), using the Gauss–Weingarten formulas, we have

$$-JA_NX + fD_XN = B(X, PN) + D_X(fN) \tag{2.20}$$

from which

$$(D_Xf)N = -B(X, PN) - JA_NX. \tag{2.21}$$

We say that if  $D_Xf = 0$  for any tangent vector field  $X$ , then the  $f$ -structure in the normal bundle is parallel.

**Lemma 2.1.** *Let  $M$  be a totally real  $n$ -dimensional submanifold of a Kaehlerian product manifold  $\overline{M}$ . If the  $f$ -structure in the normal bundle is parallel, then we have*

$$A_N = 0 \quad \text{for } N \in N_x(M). \tag{2.22}$$

**Proof.** If  $N \in N_x(M)$ , then we have  $PN = 0$ . Thus, by the assumption and (2.21) we have (2.22).  $\square$

**Remark.** We can define a frame  $e_1^*, \dots, e_n^*$  for  $JT_x(M)$  and a frame  $e_{n+1}, \dots, e_{n+p}, e_{(n+1)^*}, \dots, e_{(n+p)^*}$  for  $N_x(M)$ . Therefore if the  $f$ -structure in the normal bundle is parallel, then we have

$$A_\lambda = 0, \quad \text{i.e., } h^\lambda_{ij} = 0. \tag{2.23}$$

### 3. Integral formulas

Let  $M$  be a totally real submanifold of real dimension  $n$  of Kaehlerian product manifold  $\overline{M}(c)$  of complex dimension  $n + p$  and of constant holomorphic sectional curvature  $c$ . We prove the following lemma for later use.

**Lemma 3.1.** *Let  $M$  be a totally real submanifold of Kaehlerian product manifold  $\overline{M}(c) = \overline{M}_1^n(c_1) \times \overline{M}_2^p(c_2)$ . Then, we have*

$$\begin{aligned} \sum_{a,i,j} h^a_{ij} \Delta h^a_{ij} &= \sum_{a,i,j,k} h^a_{ij} h^a_{kkij} \\ &+ \frac{1}{16} (c_1 + c_2) \sum_a \{ [n + 9 + 15(TrFJ)^2 + 6(TrF)^2] TrA_a^2 \\ &- (3 + (TrF)^2 + (TrFJ)^2) (TrA_a)^2 \} \\ &+ \frac{1}{16} (c_1 - c_2) \sum_a \{ (n + 1) (TrF) TrA_a^2 - 2(TrF) (TrA_a)^2 \} \\ &+ \frac{1}{16} (c_1 + c_2) \sum_t \{ [4(TrF)^2 - 2] TrA_t^2 - (1 + (TrF)^2) (TrA_t)^2 \} \\ &+ \frac{1}{16} (c_1 - c_2) \sum_t \{ [2(TrF) (TrA_t^2) - 2(TrF) (TrA_t)^2] \\ &+ \frac{1}{16} (c_1 - c_2) \sum (TrJF) g(Je_b, e_a) h^a_{ii} h^b_{kk} \\ &- \frac{7}{16} (c_1 - c_2) \sum (TrJF) g(Je_b, e_a) h^b_{jj} h^a_{ij} \\ &+ \sum_{a,b} \{ Tr(A_a A_b - A_b A_a)^2 - [Tr(A_a A_b)]^2 - TrA_b Tr(A_a A_b A_a) \} \end{aligned} \tag{3.1}$$

where we have put  $A_t = A_{t^*}$ .

**Proof.** By a straightforward computation, we have [12]

$$\begin{aligned} \sum_{a,i,j} h^a_{ij} \Delta h^a_{ij} &= \sum_{a,i,j,k} (h^a_{ij} h^a_{kkij} - K^a_{ijb} h^b_{ij} h^a_{kk} + 4K^a_{bki} h^b_{jk} h^a_{ij} - K^a_{kbi} h^a_{ij} h^b_{ij} \\ &+ 2K^l_{kik} h^a_{ij} h^a_{ij} + 2K^l_{ijk} h^a_{ik} h^a_{ij}) \\ &- \sum_{a,b,i,j,k,l} [(h^a_{ik} h^b_{jk} - h^a_{jk} h^b_{ik}) (h^a_{il} h^b_{jl} - h^a_{jl} h^b_{il}) \\ &+ h^a_{ij} h^a_{kl} h^b_{ij} h^b_{kl} - h^a_{ji} h^a_{kl} h^b_{kj} h^b_{il}]. \end{aligned}$$

Then in light of above equation, using (2.1) we have our assertion.  $\square$

Now, we use Lemma 2.1 and Eq. (3.1) to obtain the following result.

**Lemma 3.2.** *Let  $M$  be a totally real submanifold of Kaehlerian product manifold  $\overline{M}(c) = \overline{M}_1^n(c_1) \times \overline{M}_2^p(c_2)$ . If the  $f$ -structure in the normal bundle is parallel, then we have*

$$\begin{aligned}
\sum_{a,i,j} h_{ij}^a \Delta h_{ij}^a &= \sum_{a,i,j,k} h_{ij}^a h_{kkij}^a \\
&+ \frac{1}{16} (c_1 + c_2) \sum_t \{ [n + 7 + 15(TrJF)^2 + 10(TrF)^2] TrA_t^2 \\
&- [4 + 2(TrF)^2 + (TrFJ)^2] (TrA_t)^2 \} \\
&+ \frac{1}{16} (c_1 - c_2) \sum_t \{ (n + 3)(TrF) TrA_t^2 - 4(TrF)(TrA_t)^2 \} \\
&+ \frac{1}{16} (c_1 - c_2) \sum (TrJF) g(Je_b, e_a) h_{ii}^a h_{kk}^b \\
&- \frac{7}{16} (c_1 - c_2) \sum (TrJF) g(Je_b, e_a) h_{ji}^b h_{ij}^a \\
&+ \sum_{t,s} \{ Tr(A_t A_s - A_s A_t)^2 \\
&- [Tr(A_t A_s)]^2 + TrA_s Tr(A_t A_s A_t) \} \tag{3.2}
\end{aligned}$$

We require the following lemma [12].

**Lemma 3.3.** *Let  $A$  and  $B$  be symmetric  $(n, n)$ -matrices. Then*

$$-Tr(AB - BA)^2 \leq 2TrA^2 TrB^2$$

and the equality holds for non-zero matrices  $A$  and  $B$  if and only if  $A$  and  $B$  can be transformed by an orthogonal matrix simultaneously into scalar multiples of  $\bar{A}$  and  $\bar{B}$  respectively, where

$$\bar{A} = \left( \begin{array}{cc|c} 0 & 1 & 0 \\ 1 & 0 & 0 \\ \hline 0 & 0 & 0 \end{array} \right),$$

$$\bar{B} = \left( \begin{array}{cc|c} 1 & 0 & 0 \\ 0 & -1 & 0 \\ \hline 0 & 0 & 0 \end{array} \right),$$

Moreover, if  $A_1, A_2, A_3$  are three symmetric  $(n, n)$ -matrices such that

$$-Tr(A_a A_b - A_b A_a)^2 = 2TrA_a^2 TrB_b^2, \quad 1 \leq a, b \leq 3, \quad a \neq b$$

then at least one of the matrices  $A_a$  must be zero.

We next put  $S_{ab} = \sum_{i,j} h_{ij}^a h_{ij}^b = TrA_a A_b$ ,  $S_a = S_{aa}$ ,  $S = \sum_a S_a$ , so that  $S_{ab}$  is a symmetric  $(n, n)$ -matrix and can be assumed to be diagonal for a suitable frame.  $S$  is the square of the length of the second fundamental form. When the  $f$ -structure in the normal bundle is parallel, using these notations, we can rewrite (3.2) in the following form:

$$\begin{aligned}
\sum_{a,i,j} h_{ij}^a \Delta h_{ij}^a &= \sum_{a,i,j,k} h_{ij}^a h_{kkij}^a + \frac{1}{16} (c_1 + c_2) [n + 7 + 15(TrJF)^2 \\
&+ 10(TrF)^2] S + \frac{1}{16} (c_1 - c_2) (n + 3)(TrF) S \\
&- \sum_t S_t^2 + \sum_{t,s} \{ Tr(A_t A_s - A_s A_t)^2 \\
&- \frac{1}{16} (c_1 + c_2) \sum_t [4 + 2(TrF)^2 + (TrFJ)^2] (TrA_t)^2 \\
&- \frac{1}{16} (c_1 - c_2) \sum_t 4(TrF)(TrA_t)^2 + \sum_{t,s} TrA_s Tr(A_t A_s A_t) \\
&+ \frac{1}{16} (c_1 - c_2) \sum (TrJF) g(Je_b, e_a) h_{ii}^a h_{kk}^b \\
&- \frac{7}{16} (c_1 - c_2) \sum (TrJF) g(Je_b, e_a) h_{ji}^b h_{ij}^a \} \tag{3.3}
\end{aligned}$$

Now, we prove the following theorem.

**Theorem 3.4.** *Let  $M$  be a compact orientable totally real submanifold of Kaehlerian product manifold  $\bar{M}(c) = \bar{M}_1^n(c_1) \times \bar{M}_2^p(c_2)$ . If the  $f$ -structure in the normal bundle is parallel, then we have*

$$\begin{aligned}
\int_M \left[ W - \sum_a (TrA_a) \Delta (TrA_a) \right] * 1 &\geq \int_M \sum_{a,i,j,k} (h_{ijk}^a)^2 * 1 \\
&\geq 0 \tag{3.4}
\end{aligned}$$

where

$$\begin{aligned}
W &= \left( 2 - \frac{1}{n} \right) S^2 - \frac{1}{16} (c_1 + c_2) [n + 7 \\
&+ 15(TrJF)^2 + 10(TrF)^2] S - \frac{1}{16} (c_1 - c_2) (n + 3)(TrF) S \\
&+ \frac{1}{16} (c_1 + c_2) \sum_t [4 + 2(TrF)^2 + (TrFJ)^2] \sum_t (TrA_t)^2 \\
&+ \frac{1}{16} (c_1 - c_2) \sum_t 4(TrF)(TrA_t)^2 - \sum_t TrA_s Tr(A_t A_s A_t) \\
&- \frac{1}{16} (c_1 - c_2) \sum (TrJF) g(Je_b, e_a) h_{ii}^a h_{kk}^b \\
&+ \frac{7}{16} (c_1 - c_2) \sum (TrJF) g(Je_b, e_a) h_{ji}^b h_{ij}^a.
\end{aligned}$$

**Proof.** Taking into account Lemma 3.3, we have

$$\begin{aligned}
&- \sum_{t,s} Tr(A_t A_s - A_s A_t)^2 + \sum_t S_t^2 \\
&- W_1 S - W_2 S \leq 2 \sum_{t \neq s} S_t S_s + \sum_t S_t^2 - W_1 S - W_2 S \\
&= \left[ \left( 2 - \frac{1}{n} \right) S - W_1 - W_2 \right] S - \frac{1}{n} \sum_{t > s} (S_t - S_s)^2 \tag{3.5}
\end{aligned}$$

where  $W_1 = \frac{1}{16} (c_1 + c_2) [n + 7 + 15(TrJF)^2 + 10(TrF)^2]$ ,  $W_2 = \frac{1}{16} (c_1 - c_2) (n + 3)(TrF)$ . In view of (3.3) and (3.5), we have

$$- \sum_{a,i,j} h_{ij}^a \Delta h_{ij}^a \leq W - \sum_{a,i,j,k} h_{ijk}^a \tag{3.6}$$

where

$$\begin{aligned}
W &= \left[ \left( 2 - \frac{1}{n} \right) S - W_1 - W_2 \right] S + \frac{1}{16} (c_1 + c_2) \sum_t [4 + 2(TrF)^2 \\
&+ (TrFJ)^2] \sum_t (TrA_t)^2 + \frac{1}{16} (c_1 - c_2) \sum_t 4(TrF)(TrA_t)^2 \\
&- \sum_t TrA_s Tr(A_t A_s A_t) \\
&- \frac{1}{16} (c_1 - c_2) \sum (TrJF) g(Je_b, e_a) h_{ii}^a h_{kk}^b \\
&+ \frac{7}{16} (c_1 - c_2) \sum (TrJF) g(Je_b, e_a) h_{ji}^b h_{ij}^a.
\end{aligned}$$

Let us assume that  $M$  is compact and orientable, then we have the following integral formulas [6]:

$$\int_M \sum_{a,i,j,k} (h_{ijk}^a)^2 * 1 = - \int_M h_{ij}^a \Delta h_{ij}^a * 1,$$

$$\int_M \sum_{a,i,j,k} h_{ij}^a h_{kij}^a * 1 = \int_M \sum_a (Tr A_a) \Delta(Tr A_a) * 1.$$

Using inequality (3.6) and above integral formulas we have Eq. (3.4) and this completes the proof of the Theorem.  $\square$

**Theorem 3.5.** *Let  $M$  be a compact orientable totally real minimal submanifold of Kaehlerian product manifold  $\overline{M}(c)$ . If the  $f$ -structure in the normal bundle is parallel, then we have*

$$\begin{aligned} \int_M \left[ \left( 2 - \frac{1}{n} \right) S - \frac{1}{16} (c_1 + c_2) \{ n + 7 + 15(Tr JF)^2 + 10(Tr F)^2 \} \right. \\ \left. - \frac{1}{16} (c_1 - c_2) (n + 3) (Tr F) \right] S * 1 - \int_M \sum_{a,i,j,k} (h_{ijk}^a)^2 * 1 \\ \geq 0 \end{aligned} \quad (3.7)$$

We finish this section by stating the following corollary.

**Corollary 3.6.** *Let  $M$  be a compact orientable totally real minimal submanifold of Kaehlerian product manifold  $\overline{M}(c)$  of complex dimension  $n + p$ . If the  $f$ -structure in the normal bundle is parallel and if*

$$\begin{aligned} S < \frac{1}{16} \frac{n}{(2n-1)} (c_1 + c_2) \{ n + 7 + 15(Tr JF)^2 + 10(Tr F)^2 \} \\ + \frac{1}{16} \frac{n}{(2n-1)} (c_1 - c_2) (n + 3) (Tr F), \end{aligned}$$

then  $M$  is totally geodesic.

#### Acknowledgments

I am thankful to Department of Science and Technology, Government of India, for its financial assistance provided through

Inspire Fellowship No. DST/INSPIRE Fellowship/2009/[xxv] to carry out this research work. I also express my sincere thanks to Reviewers for their valuable suggestions that help me improve this paper.

#### References

- [1] K. Yano, M. Kon, Totally real submanifolds of complex space form I, *Tohoku Math. J.* 28 (1976) 215–225.
- [2] K. Yano, M. Kon, Totally real submanifolds of complex space form II, *Kodai Math. Sem. Rep.* 27 (1976) 385–399.
- [3] C.S. Houch, in: Some totally real minimal surfaces in  $C^2$ , *Proc. Am. Math. Soc.* 40 (1973) 240–244.
- [4] S.T. Yau, Submanifolds with constant mean curvature I, *Am. J. Math.* 96 (1974) 346–366.
- [5] B.Y. Chen, K. Ogiue, On totally real manifolds, *Trans. Am. Math. Soc.* 193 (1974) 257–266.
- [6] M. Kon, Totally real submanifolds in Kaehler manifold, *J. Differential. Geom.* 11 (2) (1976) 251–257.
- [7] G.D. Ludden, M. Okumura, K. Yano, Totally real submanifolds of complex manifolds, *Atti Accad. Naz. Lincei Rend. Cl. Sci. Fiz. Mat. Natur.* 58 (1975) 346–353.
- [8] G.D. Ludden, M. Okumura, K. Yano, A totally real surface in  $CP^2$  that is not totally geodesic, *Proc. Am. Math. Soc.* 53 (1975) 186–190.
- [9] K. Yano, Differential geometry of totally real submanifolds, *Topics Differential. Geom.* 11 (1976) 351–359.
- [10] K. Yano, M. Kon, Structures on manifolds, in: *Series in Pure Mathematics*, vol. 3, World Scientific, Singapore, 1984.
- [11] K. Yano, On a structure defined by a tensor field  $f$  of type  $(1, 1)$   $f^2 + f = 0$ , *Tensor*, NS 14 (1963) 99–109.
- [12] S.S. Chern, M. Do Carmo, S. Kobayashi, Minimal submanifolds of a sphere with second fundamental form of constant length, *Functional Analysis and Related Fields*, Springer-Verlag, 1970, pp. 57–75.