




Egyptian Mathematical Society
Journal of the Egyptian Mathematical Society

www.etms-eg.org
www.elsevier.com/locate/joems



Original Article

Some studies of the interaction between N-two level atoms and three level atom



D.A.M. Abo-Kahla ^{a,*}, M.M.A. Ahmed ^b

^a Department of Mathematics, Faculty of Education, Ain Shams University, Roxy, Cairo, Egypt

^b Department of Mathematics, Faculty of Science, Al-Azhar University, Nasr City 11884, Cairo, Egypt

Received 6 June 2015; revised 19 September 2015; accepted 29 October 2015

Available online 10 December 2015

Keywords

Purity;
Atomic inversion;
Systems of N-two level atoms

Abstract In this paper, we present the analytical solution for the model that describes the interaction between a three level atom and two systems of N-two level atoms. The effect of the quantum numbers on the atomic inversion and the purity, for some special cases of the initial states, are investigated. We observe that the atomic inversion and the purity change remarkably by the change of the quantum numbers.

2010 Mathematics Subject Classification: 78-81

Copyright 2015, Egyptian Mathematical Society. Production and hosting by Elsevier B.V.
This is an open access article under the CC BY-NC-ND license
(<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

The use of statistical mechanics is fundamental to the concepts of quantum optics: light is described in terms of field operators for creation and annihilation of photons [1–7]. The development of quantum optics started in the 50s and 60s, motivated by the need to understand photodetection, statistics and coherence of light within the framework of quantum theory. In the late 50s,

the seminal Hanbury-Brown and Twiss (HBT) experiment [8,9] demonstrated the bunching of photons emitted by a classical chaotic source and stimulated the development of the quantum theory of optical coherences by Glauber [10–12]. An important milestone in quantum optics was the demonstration of the non-classical character of fluorescence radiation emitted by a single atom and, later, the demonstration of photons sources showing photon antibunching [13–15]. In recent years the quantum optics of attenuating [16–22] and amplifying [23–27] dielectric media was developed, where optical modes are described as open quantum systems. Features of quantum optics are mainly based on three different types of interaction, namely, field–field, atom–atom, atom–field interaction. These interactions have been extensively considered in a huge number of papers; see for example Refs. [28–44]. Each one of these interactions represents a certain type of physical phenomena [45–50]. In fact these types of interactions have a strong relation between them so that one

* Corresponding author: Tel.: +20 222440133.

E-mail address: doaa_abukahla@gmail.com (D.A.M. Abo-Kahla).
Peer review under responsibility of Egyptian Mathematical Society.



Production and hosting by Elsevier

can transform one type to another [51]. This particular kind of the transformation would enable us to transform field–field interaction into atom–field interaction or atom–atom interaction. This depends on the nature of which kind of the Hamiltonian we use to carry out the transformation. The studying of any kind of these interaction would surely lead to more progress in the field of quantum optics and consequently it may open the door to observe or to report a new phenomenon in this field [52]. In the present communication we are concerned with the type of atom–atom (spin–spin) interaction, the interaction between a three level atom and two systems of N-two level atoms. The time evolution of dynamical systems has attracted considerable attention over the past several decades because of its various applications [53]. An important aspect in this regard is the atomic inversion and the purity. The atomic population inversion can be considered as one of the simplest important quantities, it is defined as “the difference between the probabilities of finding the atom in its excited state and in its ground state”. The purity can range between zero, corresponding to a completely pure state, and $(1 - \frac{1}{d})$, corresponding to a completely mixed state, (here, d is the dimension of the density matrix). Hence, we study the atomic inversion and the purity of a three level atom coupled to two systems of N-two level atoms as an application.

This paper is organized as follows: in Section 2, we will describe the Hamiltonian of the system of interest, and obtain the explicit analytical solution of the model describing the interaction between a three level atom and two systems of N-two level atoms. Different cases are studied to demonstrate the effects due to the quantum numbers m_1, m_2 on the atomic inversion $\langle S_z \rangle$ and the purity $P_S(t)$ of the model. Conclusions are summarized at the end of the paper.

2. The model

The Hamiltonian of our model describes the interaction between a three level atom coupled to two systems of N-two level atoms. In this case the Hamiltonian of the whole system can be written in the form:

$$H = H_0 + H_{int}, \quad (1)$$

where

$$H_0 = \sum_{\alpha=1}^{\alpha=2} \omega_\alpha J_z^{(\alpha)} + \sum_{\beta=1}^{\beta=3} \Omega_\beta S_{\beta\beta}, \quad (2)$$

$$H_0 = \sum_{\gamma=1}^{\gamma=2} \omega_\gamma J_z^{(\gamma)} + \sum_{\beta=1}^{\beta=3} \Omega_\beta S_{\beta\beta}, \quad (3)$$

$$H_{int.} = \lambda_1 \left[S_{21} J_+^{(1)} + J_-^{(1)} S_{12} \right] + \lambda_2 \left[S_{32} J_+^{(2)} + J_-^{(2)} S_{23} \right], \quad (4)$$

$$J_L^{(\gamma)} = \frac{1}{2} \sum_{k_y=1}^{N_y} \sigma_L^{k_y}, L = x, y, z, \quad (5)$$

$$\Omega_1 > \Omega_2 > \Omega_3, \quad (6)$$

$$J_\pm^{(\gamma)} = J_x^{(\gamma)} \pm i J_y^{(\gamma)}, \quad (7)$$

where $\omega_\gamma, \gamma = 1, 2$ are the strength of the field (the two systems of N-two level atoms). The operators S_{ij} satisfy the commutation relation

$$[S_{kl}, S_{nm}] = S_{km}\delta_{nl} - S_{nl}\delta_{km}, \quad (8)$$

$J_\pm^{(\alpha)}$ and $J_z^{(\alpha)}$ are the collective angular momentum operators for N-two level atoms, which satisfy the relations, while $J_\pm^{(\gamma)}$ and $J_z^{(\gamma)}$ are the collective angular momentum operators for N-two level atoms, which satisfy the relations

$$[J_+^{(\gamma)}, J_-^{(\beta)}] = 2J_z^{(\gamma)}\delta_{\gamma\beta}, \quad (9)$$

$$[J_z^{(\gamma)}, J_\pm^{(\beta)}] = \pm J_\pm^{(\gamma)}\delta_{\gamma\beta}, \quad (10)$$

with the operator $\sigma_L^{k_y}$ are the usual Pauli matrices.

We define

$$|\Psi(0)\rangle = |\Psi(0)\rangle_{j,m} |\Psi(0)\rangle_s \quad (11)$$

$$|\Psi(0)\rangle_{j,m} = |j_1, m_1, j_2, m_2\rangle \quad (12)$$

$$|\Psi(0)\rangle_s = a(0)|1\rangle + b(0)|2\rangle + c(0)|3\rangle \quad (13)$$

$$|a(0)|^2 + |b(0)|^2 + |c(0)|^2 = 1 \quad (14)$$

Let

$$\begin{aligned} |\Psi(t)\rangle &= A_{m_1, m_2}(t)|1, j_1, m_1, j_2, m_2\rangle \\ &\quad + B_{m_1, m_2}(t)|2, j_1, m_1 + 1, j_2, m_2\rangle \\ &\quad + C_{m_1, m_2}(t)|3, j_1, m_1 + 1, j_2, m_2 + 1\rangle. \end{aligned} \quad (15)$$

From Schrödinger equation

$$i\frac{\partial}{\partial t}|\Psi(t)\rangle = H|\Psi(t)\rangle, \quad (16)$$

we get from Eqs. (1) and (15)

$$i\frac{dA_{m_1, m_2}(t)}{dt} = \alpha_1 A_{m_1, m_2}(t) + \Lambda_1(m_1)B_{m_1, m_2}(t), \quad (17)$$

$$\begin{aligned} i\frac{dB_{m_1, m_2}(t)}{dt} &= \alpha_2 B_{m_1, m_2}(t) + \Lambda_1(m_1)A_{m_1, m_2}(t) \\ &\quad + \Lambda_2(m_2)C_{m_1, m_2}(t), \end{aligned} \quad (18)$$

$$i\frac{dC_{m_1, m_2}(t)}{dt} = \alpha_3 C_{m_1, m_2}(t) + \Lambda_2(m_2)B_{m_1, m_2}(t), \quad (19)$$

where

$$\alpha_1 = (\omega_1 m_1 + \omega_2 m_2 + \Omega_1), \quad (20)$$

$$\alpha_2 = \omega_1(m_1 + 1) + \omega_2 m_2 + \Omega_2, \quad (21)$$

$$\alpha_3 = \omega_1(m_1 + 1) + \omega_2(m_2 + 1) + \Omega_3, \quad (22)$$

$$\Lambda_k = \lambda_k \sqrt{(j_k - m_k)(j_k + m_k + 1)} \quad k = 1, 2, \quad (23)$$

where λ_k the coupling parameters between spins.

Define

$$\begin{aligned} A_{m_1, m_2}(t) &= a(t)e^{-i\alpha_1 t}, B_{m_1, m_2}(t) = b(t)e^{-i\alpha_2 t}, \\ C_{m_1, m_2}(t) &= c(t)e^{-i\alpha_3 t} \end{aligned} \quad (24)$$

by substituting from Eq. (24) in Eqs. (17)–(19) we get the following equations:

$$i\frac{da(t)}{dt}e^{-i\alpha_1 t} = \Lambda_1(m_1)b(t)e^{-i\alpha_2 t}, \quad (25)$$




Fig. 1 Scheme of the interaction between a three level atom coupled to two systems of N-two level atoms.

$$i \frac{db(t)}{dt} e^{-i\alpha_2 t} = \Lambda_1(m_1)a(t)e^{-i\alpha_1 t} + \Lambda_2(m_2)c(t)e^{-i\alpha_3 t}, \quad (26)$$

$$i \frac{dc(t)}{dt} e^{-i\alpha_3 t} = \Lambda_2(m_2)b(t)e^{-i\alpha_2 t}, \quad (27)$$

from Fig. 1 we set

$$\Delta = [\omega_1 - (\Omega_1 - \Omega_2)] = [(\Omega_2 - \Omega_3) - \omega_2]. \quad (28)$$

So, we can write Eqs. (25)–(27) as the following:

$$i \frac{da(t)}{dt} = \Lambda_1 b(t) e^{-i\Delta t}, \quad (29)$$

$$i \frac{db(t)}{dt} = \Lambda_1 a(t) e^{i\Delta t} + \Lambda_2 c(t) e^{i\Delta t}, \quad (30)$$

$$i \frac{dc(t)}{dt} = \Lambda_2 b(t) e^{-i\Delta t}. \quad (31)$$

We solve Eqs. (29)–(31) analytically, we get:

$$b(t) = e^{\frac{i\Delta}{2}t} (q_1 e^{i\mu t} + q_2 e^{-i\mu t}), \quad (32)$$

where

$$\mu = \sqrt{\Delta^2 + 4(\Lambda_1^2 + \Lambda_2^2)}, \quad (33)$$

$$q_1 = \frac{\Lambda_1 a(0) + \Lambda_2 c(0) + (\frac{\Delta}{2} + \mu)b(0)}{2\mu}, \quad (34)$$

$$q_2 = \frac{-\Lambda_1 a(0) - \Lambda_2 c(0) - (\frac{\Delta}{2} - \mu)b(0)}{2\mu}, \quad (35)$$

similarly

$$a(t) = -\Lambda_1 e^{-\frac{i\Delta}{2}t} \left(\frac{q_1 e^{i\mu t}}{\mu - \frac{\Delta}{2}} - \frac{q_2 e^{-i\mu t}}{\mu + \frac{\Delta}{2}} \right) + q_3, \quad (36)$$




Fig. 2 Figures of the case in which $\Delta = 0$, $\lambda_1 = \lambda_2 = 1$ and $j_1 = 30$, $j_2 = 20$ and the initial state is $|\Psi(0)\rangle_s = |1\rangle$ where the dashed, bold solid, gray solid curves correspond, respectively, to $m_1 = m_2 = 1, 18, 20$.

$$q_3 = a(0) + \Lambda_1 \left(\frac{q_1}{\mu - \frac{\Delta}{2}} - \frac{q_2}{\mu + \frac{\Delta}{2}} \right) \quad (37)$$

and

$$c(t) = -\Lambda_2 e^{-\frac{i\Delta}{2}t} \left(\frac{q_1 e^{i\mu t}}{\mu - \frac{\Delta}{2}} - \frac{q_2 e^{-i\mu t}}{\mu + \frac{\Delta}{2}} \right) + q_4, \quad (38)$$

$$q_4 = c(0) + \Lambda_2 \left(\frac{q_1}{\mu - \frac{\Delta}{2}} - \frac{q_2}{\mu + \frac{\Delta}{2}} \right) \quad (39)$$

So from Eq. (24) we get $A_{m_1, m_2}(t)$, $B_{m_1, m_2}(t)$ and $C_{m_1, m_2}(t)$.

3. The atomic inversion and the purity

The atomic population inversion is given by

$$\langle S_z \rangle = |A_{m_1, m_2}(t)|^2 - |B_{m_1, m_2}(t)|^2 - |C_{m_1, m_2}(t)|^2. \quad (40)$$

The evolution of the purity $P_S(t)$ is given by

$$P_S(t) = 1 - Tr_S(\rho_S^2(t)) \quad (41)$$

where $\rho_S(t)$ is the reduced density matrix of the system which is defined by $\rho_S(t) = Tr_F \rho(t)$.

From Eq. (41) the purity can range between zero, corresponding to a completely pure state, and $(1 - \frac{1}{d})$, corresponding to a completely mixed state. (Here, d is the dimension of the density matrix.)

Based on the analytical solution in the previous section, we shall examine the evolution in time of the atomic inversion and the purity. In the numerical results we consider ($\Delta = 0$, $\lambda_1 = \lambda_2 = 1$ and $j_1 = 30$, $j_2 = 20$). We investigate the effect of the quantum numbers m_1, m_2 on the atomic inversion and the purity. In Fig. 2, the initial state is $|\Psi(0)\rangle_s = |1\rangle$. The atomic inversion and the purity have regular and periodic oscillations, when the quantum numbers $m_1 = m_2 = 1$, we find generating small oscillation between each two oscillations. When $m_1(m_2)$ increases, the phase of the generating small oscillation increases gradually until it reaches its maximum value. This means that the number of oscillations increases. In Fig. 3, the initial state is $|\Psi(0)\rangle_s = \frac{1}{\sqrt{2}}|1\rangle + \frac{1}{\sqrt{2}}|2\rangle$, when the quantum numbers $m_1 =$




Fig. 3 Figures of the case in which $\Delta = 0$, $\lambda_1 = \lambda_2 = 1$ and $j_1 = 30$, $j_2 = 20$ and the initial state is $|\Psi(0)\rangle_s = \frac{1}{\sqrt{2}}|1\rangle + \frac{1}{\sqrt{2}}|2\rangle$ where the dashed, bold solid, gray solid curves correspond, respectively, to $m_1 = m_2 = 1, 18, 20$.




Fig. 4 Figure of the case in which $\Delta = 0$, $\lambda_1 = \lambda_2 = 1$ and $j_1 = 30$, $j_2 = 20$ and the initial state is $|\Psi(0)\rangle_s = \frac{1}{\sqrt{2}}|1\rangle + \frac{1}{2}|2\rangle + \frac{1}{2}|3\rangle$ where the dashed, bold solid, gray solid curves correspond, respectively, to $m_1 = m_2 = 1, 15, 18$.




Fig. 5 Figure of the case in which $\Delta = 0$, $\lambda_1 = \lambda_2 = 1$ and $j_1 = 30$, $j_2 = 20$ and the initial state is $|\Psi(0)\rangle_s = \frac{1}{\sqrt{2}}|1\rangle + \frac{1}{2}|2\rangle + \frac{1}{2}|3\rangle$ where the dashed, bold solid, gray solid curves correspond, respectively, to $m_1 = m_2 = 1, 18, 20$.

$m_2 = 1$, the atomic inversion has regular and periodic oscillations, and its minimum value is -0.9 . When the quantum numbers $m_1(m_2)$ increases, the minimum value of the atomic inversion increases also until it reaches $\langle S_z \rangle = -0.4$. However, when the quantum numbers $m_1(m_2)$ increases until it reaches the quantum numbers $m_1 = m_2 = 20$, the atomic inversion is constant curve at $\langle S_z \rangle = 0$. When $m_1 = m_2 = 1$, the purity has regular and periodic oscillations, and its minimum value is 0.3 . When the quantum numbers $m_1(m_2)$ increases, the minimum value of the purity increases also until it reaches $\langle S_z \rangle = 0.4$. However, when the quantum numbers $m_1(m_2)$ increases until it reaches the quantum numbers $m_1 = m_2 = 20$, the purity is constant curve at $P_S(t) = 0.5$. In Figs. 4 and 5 the initial state is $|\Psi(0)\rangle_s = \frac{1}{\sqrt{2}}|1\rangle + \frac{1}{2}|2\rangle + \frac{1}{2}|3\rangle$, the atomic inversion has regular and periodic oscillations, and its minimum value is -0.65 . When the quantum numbers $m_1(m_2)$ increases, the maximum value of the atomic inversion increases also until it




Fig. 6 Figures of the case in which $\Delta = 0$, $\lambda_1 = \lambda_2 = 1$ and $j_1 = 30$, $j_2 = 20$, $m_1 = 1$, $m_2 = 18$ and the initial state is $|\Psi(0)\rangle_s = |1\rangle$.

reaches $\langle S_z \rangle = 0.5$. While the increasing the quantum numbers $m_1(m_2)$ causes the purity to reach constant curve at $P_S(t) = 0.4$. This clearly shows the effect of the the quantum numbers $m_1(m_2)$.

In Fig. 6 the initial state is $|\Psi(0)\rangle_s = |1\rangle$. We investigate the effect of the quantum numbers m_1 , m_2 , $m_1 \neq m_2$, $m_1 = 1$, $m_2 = 18$, on the atomic inversion and the purity. The atomic inversion and the purity have regular and periodic oscillations. In case $m_1 = m_2 = 1$, Fig. 2, we find generating small oscillation between each two oscillations. In case $m_1 = 1$, $m_2 = 18$, Fig. 6, The phase of the generating small oscillation increases remarkably. So, if we compare between the case of $m_1 = m_2 = 18$, and the case of $m_1 = 1$, $m_2 = 18$, we find the phase of the generating small oscillation in the second case is bigger than the phase in first case.

4. Conclusion

In this paper, we solve the model that describes the interaction between a three level atom and two systems of N-two level atoms. We calculated the atomic inversion and the purity for some special cases of the initial states ($|\Psi(0)\rangle_s$) and special values of the quantum numbers $m_1(m_2)$. When the initial state is $|\Psi(0)\rangle_s = |1\rangle$, the increasing the quantum numbers $m_1(m_2)$ causes the number of oscillations of the atomic inversion and the purity to increase. When the initial state is $|\Psi(0)\rangle_s = \frac{1}{\sqrt{2}}|1\rangle + \frac{1}{\sqrt{2}}|2\rangle$, the increasing the quantum numbers $m_1(m_2)$ causes the minimum value of the atomic inversion and the purity to increase until they reach constant curves at $\langle S_z \rangle = 0$ and $P_S(t) = 0.5$. When the initial state is $|\Psi(0)\rangle_s = \frac{1}{\sqrt{2}}|1\rangle + \frac{1}{2}|2\rangle + \frac{1}{2}|3\rangle$, the increasing the quantum numbers $m_1(m_2)$ causes the maximum value of the atomic inversion to increase. While the increasing the quantum numbers $m_1(m_2)$ causes the purity to reach constant curve at $P_S(t) = 0.4$. This clearly shows the effect of the quantum numbers $m_1(m_2)$. So, we found the difference between the probabilities of finding the atom in its exited state and in its ground state depend on the values of the quantum number and the initial state of the model. Also, we can reaching to a completely pure state, or a completely mixed state by controlling of the quantum number and the initial state of the model.

Acknowledgments

We would like to express our deep gratitude to our Prof. Abdel-Shafy F. Obada, Department of Mathematics, Faculty

of Science, Al-Azhar University, for his support, care, his useful suggestions, useful discussions and for his continuous help and guidance. The authors would like to thank the reviewers for their useful comment which improved the material of the paper.

References

- [1] L. Mandel, E. Wolf, *Optical Coherence and Quantum Optics*, Cambridge, 1995.
- [2] D.F. Walls, G.J. Milburn, *Quantum Optics*, Springer, 1994.
- [3] C.W. Gardiner, P. Zoller, *Quantum Noise*, Springer, 2004.
- [4] H.M. Moya-Cessa, F. Soto-Eguibar, *Introduction to Quantum Optics*, Rinton Press, 2011.
- [5] M.O. Scully, M.S. Zubairy, *Quantum Optics*, Cambridge, 1997.
- [6] W.P. Schleich, *Quantum Optics in Phase Space*, Wiley, 2001.
- [7] M. Kira, S.W. Koch, *Semiconductor Quantum Optics*, Cambridge University Press, 2011.
- [8] R.H. Brown, R. Twiss, A test of a new type of stellar interferometer on Sirius, *Nature* 178 (1956) 1046.
- [9] R.H. Brown, R. Twiss, Correlation between photons in two coherent beams of light, *Nature* 177 (1956) 27.
- [10] R. Glauber, Photon correlations, *Phys. Rev. Lett.* 10 (1963) 84.
- [11] R. Glauber, The quantum theory of optical coherence, *Phys. Rev.* 130 (1963) 2529.
- [12] R. Glauber, Coherent and incoherent states of the radiation field, *Phys. Rev.* 131 (1963) 2766.
- [13] H.J. Kimble, M. Dagenais, L. Mandel, Photon antibunching in resonance fluorescence, *Phys. Rev. Lett.* 39 (1977) 691.
- [14] C. Hong, L. Mandel, Experimental realization of a localized one-photon state, *Phys. Rev. Lett.* 56 (1986) 58.
- [15] P. Grangier, G. Roger, A. Aspect, Experimental evidence for a photon anticorrelation effect on a beam splitter: a new light on single-photon interferences, *Europhys. Lett.* 1 (1986) 173.
- [16] B. Huttner, S.M. Barnett, Quantization of the electromagnetic field in dielectrics, *Phys. Rev. A* 46 (1992) 4306.
- [17] T. Gruner, D.-G. Welsch, Quantum-optical input–output relations for dispersive and lossy multilayer dielectric plates, *Phys. Rev.* 54 (1996) 1661.
- [18] S. Scheel, L. Knöll, D.-G. Welsch, QED commutation relations for inhomogeneous Kramers–Kronig dielectrics, *Phys. Rev. A* 58 (1998) 700.
- [19] M. Artoni, R. Loudon, Propagation of nonclassical light through an absorbing and dispersive slab, *Phys. Rev. A* 59 (1999) 2279.
- [20] L.G. Suttorp, M. Wubs, Field quantization in inhomogeneous absorptive dielectrics, *Phys. Rev. A* 70 (2004) 013816.
- [21] U. Leonhardt, T.G. Philbin, Quantum optics of spatial transformation media, *J. Opt. A: Pure Appl. Opt.* 9 (2007) S289.
- [22] T.G. Philbin, Canonical quantization of macroscopic electromagnetism, *New J. Phys.* 12 (2010) 123008.
- [23] R.J. Glauber, Theory of quantum fluctuations in optical bistability, in: *Frontiers in Quantum Optics*, Hilger, Bristol, 1986.
- [24] R. Matloob, R. Loudon, M. Artoni, S.M. Barnett, J. Jeffers, Electromagnetic field quantization in amplifying dielectrics, *Phys. Rev. A* 55 (1997) 1623.
- [25] M. Artoni, R. Loudon, Quantum theory of optical-pulse propagation through an amplifying slab, *Phys. Rev. A* 57 (1998) 622.
- [26] D.Y. Vasylyev, W. Vogel, G. Manzke, K. Henneberger, D.-G. Welsch, Nonclassicality of radiation fields propagating in complex material systems, *Phys. Status Solidi B* 246 (2009) 293.
- [27] E. Amooghorban, M. Wubs, N.A. Mortensen, F. Kheirandish, Casimir forces in multilayer magnetodielectrics with both gain and loss, *Phys. Rev. A* 84 (2011) 013806.
- [28] A.A.-O. Faisal, J. Perina, M.S. Abdalla, Statistical properties of three quantized interacting oscillators, *Phys. Scr. A* 63 (2001) 128.
- [29] M.S. Abdalla, A.A.-O. Faisal, J. Perina, Statistical properties of a solvable three-boson squeeze operator model, *Eur. Phys. J. D* 13 (2001) 423.
- [30] A.A.-O. Faisal, J. Perina, M.S. Abdalla, Generation of squeezed light in a nonlinear asymmetric directional coupler, *J. Opt. Quantum Semiclassical Opt. B* 3 (2001) 66.
- [31] A.A.-O. Faisal, J. Perina, M.S. Abdalla, Evolution of Fock states in three mixed harmonic oscillators: quantum statistics, *Int. J. Mod. Phys. B* 15 (2001) 2125.
- [32] M.S. Abdalla, A.A.-O. Faisal, J. Perina, Dynamical properties of degenerate parametric amplifier with photon-added coherent states, *Il Nuovo Cim. B* 116 (2001) 137.
- [33] M. Abdel-Aty, M.S. Abdalla, A.-S. F. Obada, Quantum information and entropy squeezing of a two-level atom with a non-linear medium, *J. Phys. A Math. Gen.* 34 (2001) 9129.
- [34] M. Abdel-Aty, M.S. Abdalla, Degree of Entanglement for the two bimodal: field interacting with two-level system, *Physica A* 307 (2002) 437.
- [35] M. Abdel-Aty, M.S. Abdalla, A.-S. F. Obada, Entropy squeezing of a two-mode multiphoton Jaynes–Cummings model in the presence of a nonlinear medium, *J. Opt. Quantum Semiclassical Opt. B* 4 (2002) 134.
- [36] M. Abdel-Aty, M.S. Abdalla, A.-S. F. Obada, Entropy and phase properties of isotropic coupled oscillators interacting with single atom: one and two-photon process, *J. Opt. Quantum Semiclassical Opt. B* 4 (2002) S133.
- [37] S. S.Hassan, M.S. Abdalla, G.A. Kader, L.-M. Hanna, Squeezing evolution with non-dissipative and dissipative SU(2) systems, *J. Opt. Quantum Semiclassical Opt. B* 4 (2002) S204.
- [38] A.A.-O. Faisal, J. Perina, J. Perinova, M.S. Abdalla, Quantum properties of a single mode dissipative amplifier against Schrödinger-Cat states, *J. Opt. Quantum Semiclassical Opt. B* 4 (2002) S153.
- [39] M.S. Abdalla, Quantum treatment of time dependent coupled oscillators, *Int. J. Mod. Phys. B* 16 (2002) 2837.
- [40] M.S. Abdalla, M. Abdel-Aty, A.-S. F. Obada, Quantum entropy of isotropic coupled oscillators interacting with single atom, *Opt. Commun.* 211 (2002) 225.
- [41] M. Abdel-Aty, M.S. Abdalla, A.S.-F. Obada, Uncertainty relation and information entropy of a time-dependent bimodel two-level system, *J. Phys. B: At. Mol. Opt. Phys.* 35 (2002) 4773.
- [42] M.S. Abdalla, M. Abdel-Aty, A.S.-F. Obada, Degree of entanglement for anisotropic coupled oscillators interacting with a single atom, *J. Opt. Quantum Semiclassical Opt. B* 4 (2002) 396.
- [43] A.A.-O. Faisal, J. Peřina, J. Peřinová, M.S. Abdalla, Evolution of cat-states in a dissipative parametric amplifier: decoherence and entanglements, *Eur. Phys. J. D* 22 (2003) 141.
- [44] M.S. Abdalla, M.M.A. Ahmed, E.M. Khalil, A.-S. F. Obada, Dynamics of an adiabatically effective two-level atom interacting with a star-like system, *Prog. Theor. Exp. Phys.* 2014 (2014) 073A02.
- [45] G.P. Agrawal, C.L. Mehta, Dynamics of parametric processes with a trilinear hamiltonian, *J. Phys. A: Math. Gen.* 7 (1974) 607.
- [46] J. Tucker, D.F. Walls, Quantum theory of parametric frequency conversion, *Ann. Phys.* 52 (1969) 1.
- [47] C.L. Tang, Spontaneous emission in the frequency up-conversion process in nonlinear optics, *Phys. Rev.* 182 (1969) 367.
- [48] J. Tucker, D.F. Walls, Quantum theory of the traveling-wave frequency converter, *Phys. Rev.* 178 (1969) 2036.
- [49] D.F. Walls, R. Barakat, Quantum-mechanical amplification and frequency conversion with a trilinear hamiltonian, *Phys. Rev. A* 1 (1970) 446.
- [50] E.T. Jayness, F.W. cumming, Comparison of quantum and semiclassical radiation theories with application to the beam maser, *Proc. IEEE* 51 (1963) 89.
- [51] M.S. Abdalla, M.M.A. Ahmed, Some statistical properties for the interaction between SU(1, 1) and SU(2) quantum systems, *J. Phys. B: At. Mol. Opt. Phys.* 43 (2010) 155503.
- [52] M.S. Abdalla, M.M.A. Ahmed, Some statistical properties for a spin-(1/2) particle coupled to two spirals, *Opt. Commun.* 285 (2012) 3578.
- [53] M. Abdel-Aty, Influence of second-order correction to Rayleigh scattering on pancharatnam phase in a three-level atom, *Mod. Phys. Lett. B* 16 (2002) 319.